

**IDEAL STRUCTURE AND SIMPLICITY
OF THE C^* -ALGEBRAS GENERATED
BY HILBERT BIMODULES**

(REVISED VERSION, JANUARY 1998)

TSUYOSHI KAJIWARA¹ CLAUDIA PINZARI² AND YASUO WATATANI¹

Department of Environmental and Mathematical Sciences,
Okayama University, Tsushima, 700, Japan

Dipartimento di Matematica,
Università di Roma Tor
Vergata, 00133 Roma, Italy

Graduate School of Mathematics,
Kyushu University, Ropponmatsu, Fukuoka, 810, Japan

ABSTRACT. Pimsner introduced the C^* -algebra $\mathcal{O}_X$ generated by a Hilbert bimodule X over a C^* -algebra $\mathcal{A}$. We look for additional conditions that X should satisfy in order to study the simplicity and, more generally, the ideal structure of $\mathcal{O}_X$ when X is finite projective. We introduce two conditions: “(I)-freeness” and “(II)-freeness”, stronger than the former, in analogy with [J. Cuntz, W. Krieger, Invent. Math. 56, 251–268] and [J. Cuntz, Invent. Math. 63, 25–40] respectively. (I)-freeness comprehend the case of the bimodules associated with an inclusion of simple C^* -algebras with finite index, real or pseudoreal bimodules with finite dimension and the case of “Cuntz–Krieger bimodules”. If X satisfies this condition the C^* -algebra $\mathcal{O}_X$ does not depend on the choice of the generators when $\mathcal{A}$ is faithfully represented.

As a consequence, if X is (I)-free and $\mathcal{A}$ is X -simple, then $\mathcal{O}_X$ is simple. In the case of Cuntz–Krieger algebras $\mathcal{O}_A$, X -simplicity corresponds to the irreducibility of the matrix A . If $\mathcal{A}$ is simple and p.i. then $\mathcal{O}_X$ is p.i., if $\mathcal{A}$ is nonnuclear then $\mathcal{O}_X$ is nonnuclear. We therefore provide examples of (purely) infinite nonnuclear simple C^* -algebras.

Furthermore if X is (II)-free, we determine the ideal structure of $\mathcal{O}_X$.

¹Supported by the Grants-in-aid for Scientific Research, The Ministry of Education, Science and Culture, Japan

²Supported by MURST, CNR–GNAFA and European Community

1. INTRODUCTION

In [Pi] Pimsner associated with every Hilbert C^* -bimodule X a C^* -algebra $\mathcal{O}_X$. The bimodule X has a natural representation in $\mathcal{O}_X$, and generates $\mathcal{O}_X$ as a C^* -algebra. As noted in [Pi], representing Hilbert C^* -bimodules in C^* -algebras is a natural generalization of the notion of Hilbert space contained in a C^* -algebra, introduced by Doplicher and Roberts (see, e.g., [DR2]).

The construction of $\mathcal{O}_X$ generalizes and unifies various examples of C^* -algebras. As noted in [Pi], $\mathcal{O}_X$ is a generalization of Cuntz–Krieger algebras $\mathcal{O}_A$ [Cu, CK], crossed product C^* -algebras $\mathcal{A} \rtimes \mathbb{Z}$ by an automorphism and crossed products by partial automorphisms introduced by Exel [Ex]. In the particular case where X is the bimodule associated with an inclusion of von Neumann algebras, the same construction had been independently considered by Katayama [Ka]. If X is a Hilbert C^* -bimodule in the sense of [BMS] $\mathcal{O}_X$, previously considered in [AEE] from the viewpoint of crossed product C^* -algebras, was introduced as a “crossed product by X ”. Furthermore, in the case where X is full and finite projective as a right $\mathcal{A}$ -module, $\mathcal{O}_X$ is the Doplicher–Roberts algebra associated to X regarded as an object of the semitensor C^* -category with objects Hilbert C^* -bimodules over $\mathcal{A}$ and arrows right adjointable maps (cf. [DPZ] or Proposition 2.5).

Examples of Hilbert C^* -bimodules are provided from inclusions of simple C^* -algebras with finite index studied in [Wa] and Izumi [I1], [I2].

Our main interest in this paper is to study the ideal structure of $\mathcal{O}_X$ when $\mathcal{A}$ is unital and X is finite projective. We shall discuss the case where $\mathcal{A}$ is σ -unital and $X_{\mathcal{A}}$ is countably generated elsewhere [P, KWP].

We look for a general criterion that generalizes and unifies the case of crossed products by properly outer automorphisms [El, OP1, Co1, Co2, CT, Ol, Ki, OP2], condition (I) and (II) for Cuntz–Krieger algebras [Cu, CK, Cu2], condition of having “determinant 1” for Doplicher and Roberts algebras [DR1, DR2, DR3], and the case of bimodules associated with inclusions of simple C^* -algebras with finite Jones index > 1 [Jo, Wa].

In §4 we formulate a condition, that we call (I)-freeness, in analogy with condition (I) in [CK]. Our main general result is Theorem 4.3. We prove that if X is (I)-free then any representation of the bimodule X in a C^* -algebra $\mathcal{D}$ that is faithful on the coefficient algebra $\mathcal{A}$ extends to a *faithful* representation of $\mathcal{O}_X$ in $\mathcal{D}$. Hence $\mathcal{O}_X$ is the unique C^* -algebra generated by a faithful representation of X in a C^* -algebra. In particular, if $\mathcal{A}$ is X -simple, $\mathcal{O}_X$ is simple.

Furthermore we introduce a stronger condition, (II)-freeness, again in analogy with property (II) of [Cu2], that allows us to determine closed ideals of $\mathcal{O}_X$ in terms of a certain class of closed ideals of $\mathcal{A}$.

The strategy of the proof of Theorem 4.3 is the following. First we identify the algebra $\mathcal{O}_X$ with the Doplicher–Roberts algebra associated with X regarded as an object of the right tensor C^* -category of Hilbert C^* -bimodules with arrows right adjointable maps (§2). Thus $\mathcal{O}_X$ carries a natural structure of $\mathbb{Z}$ -graded C^* -algebra $\bigoplus_{k \in \mathbb{Z}} \mathcal{O}_X^{(k)}$, similar to that of Cuntz–Krieger algebras and crossed product C^* -algebras by an automorphism.

We next need show that the canonical dense $*$ -subalgebra ${}^0\mathcal{O}_X$ generated by the homogeneous spaces has a unique C^* -norm and that any representation of

$\mathcal{O}_X$ that is faithful on X is faithful. To this aim it is enough to show, by a result of [DR1], that the projection of ${}^0\mathcal{O}_X$ onto $\mathcal{O}_X^{(0)}$ is continuous in any C^* -norm. This second step is present, for example, in [El, Ki] for certain crossed product C^* -algebras by outer actions of automorphism groups, in [Cu, CK] for the algebras $\mathcal{O}_n$ and $\mathcal{O}_A$, where A is a matrix that satisfies the so called condition (I), and in [DR1] for the algebras $\mathcal{O}_\rho$ associated with a unitary representation of a compact group with determinant 1. We introduce our main assumption that X is (I)-free in order to prove this second step.

It is nice to note that if X is (I)-free, the simplicity criterion that we get, that requires for $\mathcal{A}$ to be X -simple, (i.e. lack of the so called X -invariant ideals of $\mathcal{A}$), corresponds, in the case of Cuntz-Krieger algebras $\mathcal{O}_A$ to irreducibility of the matrix A .

Thirdly, in order to study the ideal structure of $\mathcal{O}_X$ we use the condition that X is (II)-free. This condition, similarly to condition (II) by Cuntz and Krieger [CK2], assures that any closed ideal $\mathcal{J}$ of $\mathcal{O}_X$ is $\mathbb{Z}$ -graded, and hence determined by a certain ideal of $\mathcal{A}$, and it guarantees that $\mathcal{O}_X/\mathcal{J}$ is still an algebra of the form $\mathcal{O}_Y$, where Y is a (I)-free Hilbert C^* -bimodule obtained as a quotient of X over a quotient of $\mathcal{A}$.

In §5 we discuss two applications of our main theorem: the case of real or pseudoreal bimodules with Jones index > 1 , that contains the special important case of bimodules associated with proper inclusions of simple C^* -algebras with finite index in the sense of [Wa], and the case of “Cuntz-Krieger” bimodules, namely bimodules for which the coefficient algebra is the direct sum of $d > 1$ simple C^* -algebras and furthermore the adjacency matrix satisfies condition (I) of [CK].

In §6 we investigate about pure infiniteness of $\mathcal{O}_X$. We prove that if X is (I)-free and $\mathcal{A}$ is simple and purely infinite then $\mathcal{O}_X$ is simple and purely infinite. There exists a relation with a classification of purely infinite simple C^* -algebras. We do not pursue this in the present paper, just see, for example, [Rø1], [Rø2] and other works of recent rapid development.

After this work was completed we received the paper [MS] by Muhly and Solel, where similar problems have been considered.

2. REPRESENTATIONS OF BIMODULE STRUCTURES

Following [Pi], a Hilbert bimodule X over a C^* -algebra $\mathcal{A}$ is a right Hilbert $\mathcal{A}$ -module (with $\mathcal{A}$ -valued inner product denoted by $(\mid)_{\mathcal{A}}$) endowed with an isometric $*$ -homomorphism $\phi : \mathcal{A} \rightarrow \mathcal{L}_{\mathcal{A}}(X)$ from $\mathcal{A}$ to the algebra of right $\mathcal{A}$ -linear adjointable maps on X (the left action of $\mathcal{A}$). In the following, we will write $X_{\mathcal{A}}$ when X is regarded just as a Hilbert $\mathcal{A}$ -module, and ${}_{\mathcal{A}}X_{\mathcal{A}}$ when it is regarded as a Hilbert $\mathcal{A}$ -bimodule.

In this section we review the various constructions of the C^* -algebra $\mathcal{O}_X$ following [Pi], [AEE], [DPZ] and [Ka]. We will assume here and in the rest of this paper, for simplicity, that the C^* -algebra $\mathcal{A}$ is unital and that the Hilbert C^* -module $X_{\mathcal{A}}$ is full and finite projective.

For $x, y \in X$, the rank one operator $\theta_{x,y} \in \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$ is defined by $\theta_{x,y}(z) = x(y|z)_{\mathcal{A}}$ for $z \in X$. We denote by $\mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}})$ the norm closure of linear combinations of rank one operators. A finite subset $\{u_1, \dots, u_n\} \subset X$ is called a *basis* for X if $x = \sum_{i=1}^n u_i(u_i|x)_{\mathcal{A}}$ for all $x \in X$. Recall that since $X_{\mathcal{A}}$ is finite

projective, it has a basis, and that every adjointable map on $X_{\mathcal{A}}$ is an element of $\mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}})$. We set $\mathcal{K} := \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$.

We fix a basis $\{u_1, \dots, u_n\}$ of $X_{\mathcal{A}}$. Let $\phi : \mathcal{A} \rightarrow \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$ be the defining left action of $\mathcal{A}$ on X . For $a \in \mathcal{A}$,

$$\phi(a)u_j = \sum_{i=1}^n u_i a_{ij}$$

where $a_{ij} := (u_i | \phi(a)u_j)_{\mathcal{A}} \in \mathcal{A}$. Then the C^* -algebra O_X is the universal C^* -algebra generated by $\mathcal{A}$ and n operators $\{S_1, \dots, S_n\}$ satisfying relations

$$S_i^* S_j = (u_i | u_j)_{\mathcal{A}}, \quad \sum_{i=1}^n S_i S_i^* = I \quad \text{and} \quad a S_j = \sum_{i=1}^n S_i a_{ij} \quad \text{for } a \in \mathcal{A},$$

for $j = 1, \dots, n$. The generators $S_i (i = 1, \dots, n)$ are partial isometries if and only if the basis satisfies $(u_i | u_j)_{\mathcal{A}} = 0$ ($i \neq j$), because the orthogonality of the basis implies that $u_i = u_i(u_i | u_i)_{\mathcal{A}}$.

The following formulation of the universality of the algebra $\mathcal{O}_X$, due to Pimsner [Pi], makes use only of the bimodule structure of X , with no need of choosing a basis of X .

A representation of the Hilbert $\mathcal{A}$ -bimodule ${}_{\mathcal{A}}X_{\mathcal{A}}$ in a C^* -algebra $\mathcal{D}$ is a pair $(V, \rho_{\mathcal{A}})$ constituted by linear map $V : X \rightarrow \mathcal{D}$ and a unital $*$ -homomorphism $\rho_{\mathcal{A}} : \mathcal{A} \rightarrow \mathcal{D}$ satisfying the following relations:

$$V_{xa} = V_x \rho_{\mathcal{A}}(a), \tag{2.1}$$

$$V_{\phi(a)x} = \rho_{\mathcal{A}}(a) V_x, \tag{2.2}$$

$$V_x^* V_y = \rho_{\mathcal{A}}((x | y)_{\mathcal{A}}) \tag{2.3}$$

for $x, y \in X, a \in \mathcal{A}$. Note that $\rho_{\mathcal{A}}$ is uniquely determined by V since X is assumed to be full. If $\{u_i, i = 1, \dots, n\}$ is a basis of X , $\sum_{i=1}^n V_{u_i} V_{u_i}^*$ does not depend on the choice of the basis, and is called the *support* of $(V, \rho_{\mathcal{A}})$.

We regard X as an imprimitivity bimodule ${}_{\mathcal{K}}X_{\mathcal{A}}$ in the sense of [Ri], with left inner product ${}_{\mathcal{K}}(x | y) = \theta_{x,y}$ for $x, y \in X$. A representation of ${}_{\mathcal{K}}X_{\mathcal{A}}$ in a C^* -algebra $\mathcal{D}$ is a triple $(V, \rho_{\mathcal{K}}, \rho_{\mathcal{A}})$ where $(V, \rho_{\mathcal{A}})$ is a representation of ${}_{\mathcal{A}}X_{\mathcal{A}}$ in $\mathcal{D}$ and $\rho_{\mathcal{K}} : \mathcal{K} \rightarrow \mathcal{D}$ is a unital $*$ -homomorphism such that

$$V_{kx} = \rho_{\mathcal{K}}(k) V_x, \tag{2.4}$$

$$V_x V_y^* = \rho_{\mathcal{K}}({}_{\mathcal{K}}(x | y)) \tag{2.5}$$

for $x, y \in X, k \in \mathcal{K}$. Note that since $\rho_{\mathcal{K}}$ is unital,

$$\sum_{i=1}^n V_{u_i} V_{u_i}^* = I, \tag{2.6}$$

for any basis $\{u_i, i = 1, \dots, n\}$ of X . Furthermore, comparing (2.2) and (2.4), $\rho_{\mathcal{K}} \circ \phi = \rho_{\mathcal{A}}$.

The C^* -algebra $\mathcal{O}_X$ is the universal C^* -algebra generated by operators S_x , $x \in X$ where $S : X \rightarrow \mathcal{O}_X$ defines a representation of the imprimitivity bimodule ${}_K X_{\mathcal{A}}$ in $\mathcal{O}_X$.

The equivalence of the above two definitions depends on the fact that any representation of the Hilbert $\mathcal{A}$ -bimodule ${}_A X_{\mathcal{A}}$ in a unital C^* -algebra $\mathcal{D}$ with support I extends to a representation of the imprimitivity bimodule ${}_K X_{\mathcal{A}}$ [Pi]. We will give a proof in Lemma 2.2 and Corollary 2.3, because it is one of the steps of the proof of the simplicity of $\mathcal{O}_X$.

[Pi] and [Ka], generalizing to Hilbert C^* -bimodules a well known construction for Hilbert spaces, provide a concrete realization of $\mathcal{O}_X$ as a quotient of the Toeplitz algebra $\mathcal{T}_X$ acting on the Fock space of X . This construction shows in particular that every Hilbert C^* -bimodule can be *faithfully* represented in some C^* -algebra.

We consider right Hilbert $\mathcal{A}$ -modules $X^{\otimes m} = X \otimes_{\mathcal{A}} \cdots \otimes_{\mathcal{A}} X$ (m -times), with the convention $X^{\otimes 0} = \mathcal{A}$ (since $X_{\mathcal{A}}$ is finite projective, the algebraic tensor products over $\mathcal{A}$ are right Hilbert $\mathcal{A}$ -modules, with no need of completion (cf. [KW1])). Let $F(X) = \bigoplus_{m=0}^{\infty} X^{\otimes m}$ be the full Fock space of X , regarded as a right Hilbert $\mathcal{A}$ -module. We consider the representation of ${}_A X_{\mathcal{A}}$ in $L_{\mathcal{A}}(F(X)_{\mathcal{A}})$ by letting $\mathcal{A}$ act in the obvious way on $F(X)$ and defining, for $x \in X$, the creation operator $T_x \in \mathcal{L}_{\mathcal{A}}(F(X)_{\mathcal{A}})$

$$\begin{aligned} T_x(x_1 \otimes \cdots \otimes x_m) &= x \otimes x_1 \otimes \cdots \otimes x_m, \\ T_x(a) &= xa \end{aligned}$$

for $x_1 \otimes \cdots \otimes x_m \in X^{\otimes m}$ and $a \in \mathcal{A}$. The C^* -subalgebra $\mathcal{T}_X$ of $\mathcal{L}_{\mathcal{A}}(F(X)_{\mathcal{A}})$ generated by $\{T_x; x \in X\}$ is called the Toeplitz algebra of X .

Let $\pi : \mathcal{L}_{\mathcal{A}}(F(X)_{\mathcal{A}}) \rightarrow \mathcal{L}_{\mathcal{A}}(F(X)_{\mathcal{A}})/\mathcal{K}_{\mathcal{A}}(F(X)_{\mathcal{A}})$ be the quotient map, and set $S_x = \pi(T_x)$, $x \in X$. Let $C^*(S_X)$ denote the C^* -algebra generated by $\{S_x; x \in X\}$. For $x = (x_1, \dots, x_r) \in X^{\times r}$, we set $S_x = S_{x_1} \cdots S_{x_r}$ and put $S_{\emptyset} = I$. Let $\mathcal{F}_{r,s}$ denote the closed subspace of $C^*(S_X)$ generated by $\{S_x S_y^*; x \in X^{\times s}, y \in X^{\times r}\}$. Let, for $T \in \mathcal{K} = \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$, $\pi_{\mathcal{K}}(T) \in \mathcal{L}_{\mathcal{A}}(F(X)_{\mathcal{A}})/\mathcal{K}_{\mathcal{A}}(F(X)_{\mathcal{A}})$ denote the image in the quotient algebra of the operator on $F(X)$ that acts trivially on $X^{\otimes 0} = \mathcal{A}$ and that on $X^{\otimes r}$, $r = 1, 2, \dots$ acts by $x_1 \otimes \cdots \otimes x_r \mapsto (Tx_1) \otimes \cdots \otimes x_r$. Since $\pi_{\mathcal{K}}(\theta_{x,y}) = S_x S_y^*$ for $x, y \in X$, we have $\pi_{\mathcal{K}}(T) \in \mathcal{F}_{1,1} \subset C^*(S_X)$ for $T \in \mathcal{K}$. Put $\pi_{\mathcal{A}}(a) = \pi_{\mathcal{K}}(\phi(a))$ for $a \in \mathcal{A}$. Then $(S, \pi_{\mathcal{K}}, \pi_{\mathcal{A}})$ is a representation of ${}_K X_{\mathcal{A}}$ in $C^*(S_X)$ with support I . It has been shown in [Pi] that $C^*(S_X)$ is the universal C^* -algebra generated by such a representation, therefore it is isomorphic to $\mathcal{O}_X$.

Let $\{u_1, \dots, u_n\}$ be a basis of X . Then the identity

$$S_x S_y^* = \sum_{i=1}^n S_x S_{u_i} S_{u_i}^* S_y^*,$$

for $x \in X^r$, $y \in X^s$, shows that $\mathcal{F}_{r,s} \subset \mathcal{F}_{r+1,s+1}$. We denote by $\mathcal{F}_{\infty}^{(k)}$ the norm closure of $\bigcup_{r=0}^{\infty} \mathcal{F}_{r,r+k}$. The unitary representation U of $\mathbb{T}$ on $F(X)$ defined by $U(t)x = t^r x$, $t \in \mathbb{T}$, $x \in X^{\otimes r}$ implements an automorphic action $\gamma : \mathbb{T} \rightarrow \text{Aut } \mathcal{O}_X$ such that $\gamma_t(S_x) = t S_x$ for $t \in \mathbb{T}$ and $x \in X$, thus making

$\mathcal{O}_X$ into a $\mathbb{Z}$ -graded C^* -algebra. The fixed point algebra $\mathcal{O}_X^\gamma$ is $\mathcal{F}_\infty^{(0)}$. We denote by m_0 the faithful conditional expectation of $\mathcal{O}_X$ onto $\mathcal{F}_\infty^{(0)}$ defined by $m_0(T) = \int_{\mathbb{T}} \gamma_t(T) dt$ for $T \in \mathcal{O}_X$.

We now recall the construction of $\mathcal{O}_X$ by [DPZ], intrinsic to the category of Hilbert C^* -bimodules. A general construction [DR3] associates functorially to each object ρ of a strict tensor C^* -category, a C^* -algebra $\mathcal{O}_\rho$. This applies, with no substantial modifications, to the case where ρ is just an object of a strict *right tensor* C^* -category $\mathcal{T}$. More specifically, $\mathcal{T}$ is a C^* -category for which the set of objects is a semigroup with unit ι , such that for any triple of objects ρ, σ, τ there are maps $L_\tau^{\rho, \sigma}$ defined on the space (ρ, σ) of arrows from ρ to σ (“tensoring on the right by 1_τ ”) that satisfy:

$$L_\tau^{\rho, \sigma} : (\rho, \sigma) \rightarrow (\rho\tau, \sigma\tau)$$

$$L_\omega^{\rho\tau, \sigma\tau} L_\tau^{\rho, \sigma} = L_{\tau\omega}^{\rho, \sigma} ,$$

$$L_\iota^{\rho, \sigma} = id .$$

These maps are assumed to satisfy natural compatibility relations with the $*$ -involution and composition of the category $\mathcal{T}$. Any strict tensor C^* -category is a strict right tensor C^* -category by $L_\tau^{\rho, \sigma}(T) = T \otimes 1_\tau$. For simplicity, we assume that $L_\tau^{\rho, \sigma}$ are isometric mappings.

Let ${}^0\mathcal{O}_\rho^{(k)}$ be the algebraic inductive limit of linear spaces (ρ^r, ρ^{r+k}) by the maps $L_\rho^{\rho^r, \rho^{r+k}}$. Then the algebraic direct sum ${}^0\mathcal{O}_\rho = \bigoplus_{k \in \mathbb{Z}} {}^0\mathcal{O}_\rho^{(k)}$ is a $\mathbb{Z}$ -graded $*$ -algebra. Let $\gamma : \mathbb{T} \rightarrow \text{Aut}({}^0\mathcal{O}_\rho)$ be the $\mathbb{T}$ -action defining the grading. The algebra $\mathcal{O}_\rho$ is defined to be the completion of ${}^0\mathcal{O}_\rho$ in the unique C^* -norm for which γ_z is isometric.

Let $\mathcal{A} \subset \mathcal{D}$ be a unital inclusion of C^* -algebras. A typical example of strict right tensor C^* -category is the category $\mathcal{H}_\mathcal{A}(\mathcal{D})$ of Hilbert $\mathcal{A}$ -bimodules contained in a C^* -algebra $\mathcal{D}$, finite projective as right $\mathcal{A}$ -modules and with support I . An object of $\mathcal{H}_\mathcal{A}(\mathcal{D})$ is a closed subspace X of $\mathcal{D}$ for which $\mathcal{A}X = X\mathcal{A} = X$ and $X^*X \subseteq \mathcal{A}$. X carries an obvious Hilbert $\mathcal{A}$ -bimodule structure. The set of arrows between two objects X and Y is the space $\mathcal{L}_\mathcal{A}(X_\mathcal{A}, Y_\mathcal{A}) = \mathcal{K}_\mathcal{A}(X_\mathcal{A}, Y_\mathcal{A})$ of right adjointable $\mathcal{A}$ -module maps, identified with the subspace YX^* of $\mathcal{D}$. The operations of composition and $*$ -involution on the arrows of $\mathcal{H}_\mathcal{A}(\mathcal{D})$ are given by the operator product and $*$ -involution inherited from $\mathcal{D}$. For any pair of objects X, Y of $\mathcal{H}_\mathcal{A}(\mathcal{D})$, XY , the linear span of operator products xy , $x \in X, y \in Y$, is a strict realization of the tensor product $X \otimes_\mathcal{A} Y$ of Hilbert $\mathcal{A}$ -bimodules. The unit object is the trivial bimodule $\mathcal{A}$. On arrows, the “tensor product with the identity arrows” is defined by $L_Z^{X, Y}(T) = T$. Note that the Hilbertian norm on X coincides with the norm inherited from $\mathcal{D}$.

Doplicher–Roberts construction applied to an object X of $\mathcal{H}_\mathcal{A}(\mathcal{D})$ provides a C^* -algebra containing X and $\mathcal{A}$. Since $X_\mathcal{A}$ is finite projective, the subspace ${}^0\mathcal{O}_\rho^{(k)}$ of $\mathcal{O}_\rho$ identifies with $\cup_r X^{r+k} X^{r*} \subset \mathcal{D}$, therefore $\mathcal{O}_\rho$ is a C^* -algebra generated by X (that in general fails to identify with the C^* -subalgebra $C^*(X)$ of $\mathcal{D}$ generated by X). It has been shown in [DPZ] that this C^* -algebra is isomorphic to $\mathcal{O}_X$. For convenience, we will give a proof of this fact. We need the following standard computation of the norm of a finite rank operator.

2.1 Lemma. *Let $\mathcal{A}$ be a C^* -algebra and $X_{\mathcal{A}}$ a right Hilbert $\mathcal{A}$ -module. For $x_1, \dots, x_n, y_1, \dots, y_n \in X$, we have*

$$\left\| \sum_{i=1}^n \theta_{x_i, y_i} \right\| = \|((x_i | x_j)_{\mathcal{A}})_{ij}^{1/2} ((y_i | y_j)_{\mathcal{A}})^{1/2}\|$$

where the right hand side is the norm on $M_n(\mathcal{A})$. In particular, if $X = \mathcal{A}$, then for $x_1, \dots, x_n, y_1, \dots, y_n \in \mathcal{A}$ we have

$$\left\| \sum_{i=1}^n x_i y_i^* \right\| = \|(x_i^* x_j)_{ij}^{1/2} (y_i^* y_j)_{ij}^{1/2}\|$$

Proof. For $x, y \in X$, we have

$$\begin{aligned} \|\theta_{x,y}\|^2 &= \|\theta_{x,y}^* \theta_{x,y}\| = \|\theta_{y(x|x)_{\mathcal{A}}, y}\| = \\ &= \|\theta_{y(x|x)_{\mathcal{A}}^{1/2}, y(x|x)_{\mathcal{A}}^{1/2}}\| = \|(y(x|x)_{\mathcal{A}}^{1/2} | y(x|x)_{\mathcal{A}}^{1/2})_{\mathcal{A}}\| = \\ &= \|(x|x)_{\mathcal{A}}^{1/2} (y|y)_{\mathcal{A}}^{1/2}\|^2. \end{aligned}$$

Consider $Y := X \oplus \dots \oplus X = X \otimes \mathbb{C}^n$ as a right $\mathcal{A} \otimes M_n(\mathbb{C})$ -module as in the proof of [KW1; Proposition 1.18]. For $x = (x_1, \dots, x_n), y = (y_1, \dots, y_n) \in Y$, we put $(x|y)_{\mathcal{A} \otimes M_n(\mathbb{C})} = ((x_i | y_j)_{\mathcal{A}})_{ij}$. Let

$$\Theta_{x,y}(z) = x(y|z)_{\mathcal{A} \otimes M_n(\mathbb{C})}$$

for $x, y, z \in Y$. Then $\Theta_{x,y} = (\sum_{i=1}^n \theta_{x_i, y_i}) \otimes I$ by the identification

$$\mathcal{L}_{\mathcal{A} \otimes M_n(\mathbb{C})}(Y_{\mathcal{A} \otimes M_n(\mathbb{C})}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}}) \otimes \mathbb{C}.$$

Therefore we have

$$\begin{aligned} \left\| \sum_{i=1}^n \theta_{x_i, y_i} \right\| &= \|\Theta_{x,y}\| = \|(x|x)_{\mathcal{A} \otimes M_n(\mathbb{C})}^{1/2} (y|y)_{\mathcal{A} \otimes M_n(\mathbb{C})}^{1/2}\| \\ &= \|((x_i | x_j)_{\mathcal{A}})_{ij}^{1/2} ((y_i | y_j)_{\mathcal{A}})^{1/2}\| \end{aligned}$$

The following Lemma shows that any representation of a right Hilbert C^* -module $X_{\mathcal{A}}$ in a C^* -algebra extends to a representation of the imprimitivity bimodule ${}_K X_{\mathcal{A}}$ in the same C^* -algebra.

Let $\mathcal{T}$ denote a full C^* -subcategory of the category $\mathcal{H}_{\mathcal{A}}$ of finite projective right Hilbert modules over $\mathcal{A}$.

2.2. Lemma. ([Pi; Lemma 3.2]) *Let $\mathcal{A}$ and $\mathcal{D}$ be unital C^* -algebras. Let be given a linear map $\pi_X : X \rightarrow \mathcal{D}$ for each object X of $\mathcal{T}$ and a unital $*$ -homomorphism $\pi_{\mathcal{A}} : \mathcal{A} \rightarrow \mathcal{D}$ such that*

$$\pi_X(xa) = \pi_X(x)\pi_{\mathcal{A}}(a) \quad \text{and} \quad \pi_X(x)^* \pi_X(y) = \pi_{\mathcal{A}}((x|y)_{\mathcal{A}})$$

for $x, y \in X$ and $a \in \mathcal{A}$. Then there exists a unique $*$ -functor $\pi : \mathcal{T} \rightarrow \mathcal{H}_{\mathcal{A}}(\mathcal{D})$ with full image that associates $\pi_X(X)$ with any object $X \in \mathcal{T}$, and such that

$$\pi(\theta_{y,x}) = \pi_Y(y)\pi_X(x)^* \quad \text{and} \quad \pi_X(kx) = \pi_{\mathcal{K}}(k)\pi_X(x)$$

for $x \in X$, $y \in Y$ and $k \in \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}, Y_{\mathcal{A}})$. Furthermore if $\pi_{\mathcal{A}}$ is one to one, then π is faithful and π_X is an isometry.

Proof. For $x_1, \dots, x_n \in X, y_1, \dots, y_n \in Y$, Lemma 2.1 with a slight modification shows that

$$\begin{aligned} \left\| \sum_{i=1}^n \theta_{x_i, y_i} \right\| &= \|((x_i|x_j)_{\mathcal{A}})_{ij}^{1/2}((y_i|y_j)_{\mathcal{A}})_{ij}^{1/2}\| \\ &\geq \|(\pi_{\mathcal{A}}((x_i|x_j)_{\mathcal{A}}))_{ij}^{1/2}(\pi_{\mathcal{A}}((y_i|y_j)_{\mathcal{A}}))_{ij}^{1/2}\| \\ &= \|(\pi_X(x_i)^* \pi_X(x_j))_{ij}^{1/2}(\pi_Y(y_i)^* \pi_Y(y_j))_{ij}^{1/2}\| \\ &= \left\| \sum_{i=1}^n \pi_X(x_i) \pi_Y(y_i)^* \right\|. \end{aligned}$$

The above estimate shows that there exists a contraction $\pi : \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}, Y_{\mathcal{A}}) \rightarrow \mathcal{D}$ such that $\pi(\theta_{y,x}) = \pi_Y(y)\pi_X(x)^*$. It is easily seen that the map $\pi : \mathcal{T} \rightarrow \mathcal{H}_{\mathcal{A}}(\mathcal{D})$ is in fact a $*$ -functor. Since

$$\begin{aligned} \pi_X(\theta_{x,y}(z)) &= \pi_X(x(y|z)_{\mathcal{A}}) = \pi_X(x)\pi_{\mathcal{A}}((y|z)_{\mathcal{A}}) \\ &= \pi_X(x)\pi_Y(y)^* \pi_X(z) = \pi(\theta_{x,y})\pi_X(z) \end{aligned}$$

we have that $\pi_X(kz) = \pi(k)\pi_X(z)$ for $k \in \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}, Y_{\mathcal{A}})$ and $z \in X$. Furthermore assume that $\pi_{\mathcal{A}}$ is one to one. Then the only inequality in the above calculation becomes an equality. Thus $\pi : \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}, Y_{\mathcal{A}}) \rightarrow \mathcal{D}$ is also one to one. Since

$$\|\pi_X(x)\|^2 = \|\pi_X(x)^* \pi_X(x)\| = \|\pi_{\mathcal{A}}((x|x)_{\mathcal{A}})\| = \|(x|x)_{\mathcal{A}}\| = \|x\|^2,$$

π is an isometry.

Let $\mathcal{T}_X$ denote the full subcategory of the category $\mathcal{H}_{\mathcal{A}}$ of finite projective right Hilbert $\mathcal{A}$ -modules, with objects $\{X^{\otimes r}, r = 0, 1, 2, \dots\}$.

2.3. Corollary. ([Pi]) *Let $\mathcal{A}$ be a unital C^* -algebra and ${}_{\mathcal{A}}X_{\mathcal{A}}$ a Hilbert $\mathcal{A}$ -bimodule, full and finite projective as a right $\mathcal{A}$ -module. Let $\phi : \mathcal{A} \rightarrow \mathcal{K} = \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$ be the defining left action of $\mathcal{A}$. Let $\mathcal{O}_X = C^*\{S_x; x \in X\}$ and let $\pi_{\mathcal{A}} : \mathcal{A} \rightarrow \mathcal{O}_X$ be the canonical embedding. Then there exist representations $(\pi_r, \pi_{\mathcal{A}})$, $r = 0, 1, 2, \dots$, of Hilbert $\mathcal{A}$ -bimodules $X^{\otimes r}$ in $\mathcal{O}_X$ with support I such that*

$$\pi_r(x_1 \otimes \dots \otimes x_r) = S_{x_1} \dots S_{x_r}.$$

Furthermore there is a faithful right tensor $*$ -functor $\pi : \mathcal{T}_X \rightarrow \mathcal{H}_{\mathcal{A}}(\mathcal{O}_X)$ such that $\pi(X^{\otimes r}) = \pi_r(X^{\otimes r})$ and

$$\pi : \theta_{x_1 \otimes \dots \otimes x_s, y_1 \otimes \dots \otimes y_r} \in \mathcal{K}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}}, X^{\otimes s}_{\mathcal{A}}) \rightarrow S_{x_1} \dots S_{x_s} S_{y_r}^* \dots S_{y_1}^* \in \mathcal{F}_{r,s}.$$

Proof. Apply Lemma 2.2 to $X^{\otimes r}$ in place of X and put $\mathcal{D} = \mathcal{O}_X$. Since

$$S_{x_r}^* \cdots S_{x_1}^* S_{y_1} \cdots S_{y_r} = \pi_{\mathcal{A}}((x_1 \otimes \cdots \otimes x_r \mid y_1 \otimes \cdots \otimes y_r)_{\mathcal{A}}),$$

there exists an isometry $\pi_r : X^{\otimes r} \rightarrow \mathcal{O}_X$ such that $\pi_r(x_1 \otimes \cdots \otimes x_r) = S_{x_1} \cdots S_{x_r}$ that satisfies

$$\pi_r(xa) = \pi_r(x)\pi_{\mathcal{A}}(a), \quad \pi_r(\phi(a)x) = \pi_{\mathcal{A}}(a)\pi_r(x)$$

and

$$\pi_r(x)^* \pi_r(y) = \pi_{\mathcal{A}}((x|y)_{\mathcal{A}}),$$

hence π_r is a faithful representation of $X^{\otimes r}$ in $\mathcal{O}_X$, clearly with support I . By Lemma 2.2, there exists a faithful *-functor $\pi : \mathcal{T}_X \rightarrow \mathcal{H}_{\mathcal{A}}(\mathcal{D})$ that satisfies the stated properties. We show that π is right tensor. Let $\{u_1, \dots, u_n\}$ be a basis of X .

$$\begin{aligned} \pi(\theta_{x_1 \otimes \cdots \otimes x_s, y_1 \otimes \cdots \otimes y_r} \otimes I) &= \pi\left(\sum_{i=1}^n \theta_{x_1 \otimes \cdots \otimes x_s \otimes u_i, y_1 \otimes \cdots \otimes y_r \otimes u_i}\right) \\ &= \sum_{i=1}^n S_{x_1} \cdots S_{x_s} S_{u_i} S_{u_i}^* S_{y_r}^* \cdots S_{y_1}^* = S_{x_1} \cdots S_{x_s} S_{y_r}^* \cdots S_{y_1}^*. \end{aligned}$$

Let $\rho = X_{\mathcal{A}}$ be a Hilbert $\mathcal{A}$ -bimodule contained in a C^* -algebra $\mathcal{D}$ with support I , regarded as an object of the strict right tensor C^* -category $\mathcal{H}_{\mathcal{A}}(\mathcal{D})$, and associate to it the Doplicher–Roberts algebra $\mathcal{O}_{\rho}$.

2.4. Lemma. *Let $\mathcal{O}_{\rho}^{(k)}$ denote the closure of ${}^0\mathcal{O}_{\rho}^{(k)}$ in $\mathcal{O}_{\rho}$. There is an isomorphism of $\mathbb{Z}$ -graded *-algebras*

$$\psi : \oplus_{k \in \mathbb{Z}} \mathcal{O}_{\rho}^{(k)} \rightarrow \oplus_{k \in \mathbb{Z}} \mathcal{F}_{\infty}^{(k)}$$

such that $\psi(x) = S_x, x \in X$. Furthermore ψ is compatible with the inclusions $\mathcal{K}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}}, X^{\otimes r+k}_{\mathcal{A}}) \hookrightarrow \mathcal{K}_{\mathcal{A}}(X^{\otimes r+1}_{\mathcal{A}}, X^{\otimes r+k+1}_{\mathcal{A}})$ in $\mathcal{O}_{\rho}^{(k)}$ and $\mathcal{F}_{r,r+k} \subset \mathcal{F}_{r+1,r+k+1}$ in $\mathcal{F}_{\infty}^{(k)}$.

Proof. Since the *-functor π defined in Corollary 2.3 is compatible with right tensorization with the identity arrows, we may define a *-isomorphism of $\mathbb{Z}$ -graded *-algebras $\psi : \oplus_{k \in \mathbb{Z}} \mathcal{O}_{\rho}^{(k)} \rightarrow \oplus_{k \in \mathbb{Z}} \mathcal{F}_{\infty}^{(k)}$ by letting

$$\psi|_{\mathcal{K}(X^{\otimes r}_{\mathcal{A}}, X^{\otimes s}_{\mathcal{A}})} = \pi.$$

As a consequence of the previous lemma we deduce the following result, proved also in [DPZ].

2.5. Proposition. *The isomorphism $\psi : \oplus_{k \in \mathbb{Z}} \mathcal{O}_{\rho}^{(k)} \rightarrow \oplus_{k \in \mathbb{Z}} \mathcal{F}_{\infty}^{(k)}$ of $\mathbb{Z}$ -graded *-algebras defined in the previous lemma extends to an isomorphism $\psi : \mathcal{O}_{\rho} \rightarrow \mathcal{O}_X$ of C^* -algebras.*

Proof. In view of Lemma 2.4 we need to show that the norm of $\oplus_{k \in \mathbb{Z}} \mathcal{F}_{\infty}^{(k)}$ inherited from $\mathcal{L}_{\mathcal{A}}(F(X)_{\mathcal{A}})/\mathcal{K}_{\mathcal{A}}(F(X)_{\mathcal{A}})$ makes the canonical action of $\mathbb{T}$ continuous. But this is clear since this action is implemented by the quotient image of a unitary representation of $\mathbb{T}$ on $F(X)_{\mathcal{A}}$.

In the following we shall identify $\mathcal{O}_X$ with the algebra $\mathcal{O}_\rho$, omitting the isomorphism ψ . In particular, X and $\mathcal{A}$, as well as the spaces $\mathcal{K}_\mathcal{A}(X^{\otimes r}_\mathcal{A}, X^{\otimes r}_\mathcal{A})$, will be regarded as closed subspaces of $\mathcal{O}_X$.

The existence of the following sequence of conditional expectations will be useful later.

2.6. Lemma. *Consider the same situation as in Corollary 2.3. Assume furthermore that X is a Hilbert $\mathcal{A}$ -bimodule of finite type in the sense of [KW1] with left inner product ${}_\mathcal{A}(\cdot|\cdot)$. Then there exist conditional expectations $E_r : \mathcal{O}_X^{(0)} \rightarrow \mathcal{K}_\mathcal{A}(X^{\otimes r}_\mathcal{A})$ (for $r = 0, 1, 2, \dots$) such that $T = \lim_{r \rightarrow \infty} E_r(T)$ (in the norm topology) for $T \in \mathcal{O}_X^{(0)}$. Moreover there exists a conditional expectation $E_\mathcal{A}^{\mathcal{O}_X} : \mathcal{O}_X \rightarrow \mathcal{A}$*

Proof. As in [KW1, Lemma 3.23, 3.24], there exists a conditional expectation $E_r^{r+k} : \mathcal{K}_\mathcal{A}(X^{\otimes r}_\mathcal{A} \otimes_\mathcal{A} X^{\otimes k}_\mathcal{A}) \rightarrow \mathcal{K}_\mathcal{A}(X^{\otimes r}_\mathcal{A})$ such that

$$E_r^{r+k}(\theta_{x_1 \otimes y_1, x_2 \otimes y_2}) = (r - \text{Ind}[X^{\otimes k}])^{-1} \theta_{x_1 \mathcal{A}(y_1|y_2), x_2}$$

for $x_1, x_2 \in X^{\otimes r}$ and $y_1, y_2 \in X^{\otimes k}$. We also have a conditional expectation $G : \mathcal{K}_\mathcal{A}(X_\mathcal{A}) \rightarrow \mathcal{A}$ such that $G(\theta_{x,y}) = (r - \text{Ind}[X])^{-1} {}_\mathcal{A}(x|y)$ for $x, y \in X$ by [KW1; Lemma 1.26]. Then we can easily construct the desired conditional expectations $E_r : \mathcal{O}_X^{(0)} \rightarrow \mathcal{K}_\mathcal{A}(X^{\otimes r}_\mathcal{A})$. Put $E_0 = G \circ E_1$ and $E_\mathcal{A}^{\mathcal{O}_X} = E_0 \circ m_0$, where m_0 is the conditional expectation of $\mathcal{O}_X$ onto $\mathcal{O}_X^{(0)}$ obtained overaging over the action of $\mathbb{T}$.

Lemma 2.6 shows that, if X is a Hilbert $\mathcal{A}$ -bimodule of finite type, the relative commutant algebra $\mathcal{A}' \cap \mathcal{O}_X^{(0)}$ is approximated by the subalgebras $\cup_{r=1}^\infty \mathcal{A}' \cap \mathcal{K}_\mathcal{A}(X^{\otimes r}_\mathcal{A})$ (cf [KW1]).

3. CANONICAL COMPLETELY POSITIVE MAPS

Let $u_1, \dots, u_n$ be a finite basis of $X = X_\mathcal{A}$. We define a completely positive map $\sigma : \mathcal{O}_X \rightarrow \mathcal{O}_X$ by $\sigma(T) = \sum_{i=1}^n u_i T u_i^*$. We note that σ does depend on the choice of the basis, and that it is not a $*$ -endomorphism in general. However,

3.1. Lemma. *The restriction of σ to $\mathcal{A}' \cap \mathcal{O}_X$ is a unital $*$ -monomorphism that does not depend on the choice of the basis of $X_\mathcal{A}$.*

Proof. For $T_1, T_2 \in \mathcal{A}' \cap \mathcal{O}_X$,

$$\begin{aligned} \sigma(T_1)\sigma(T_2) &= \sum_i u_i T_1 u_i^* \sum_j u_j T_2 u_j^* \\ &= \sum_i \sum_j u_i T_1 (u_i|u_j)_\mathcal{A} T_2 u_j^* = \sum_i \sum_j u_i (u_i|u_j)_\mathcal{A} T_1 T_2 u_j^* \\ &= \sum_j u_j T_1 T_2 u_j^* = \sigma(T_1 T_2). \end{aligned}$$

Suppose that $\sigma(T) = 0$ for $T \in \mathcal{A}' \cap \mathcal{O}_X$. Then $\sigma(TT^*) = \sum_i u_i T T^* u_i^* = 0$. Hence $u_i T = 0$. Then $u_i a T = u_i a T = u_i T a = 0$ for $a \in \mathcal{A}$. Thus $xT = 0$ for all $x \in X$. Therefore $(y|x)_\mathcal{A} T = y^* x T = 0$. Since $X_\mathcal{A}$ is full, $T = 0$. Thus

σ is isometric on $\mathcal{A}' \cap \mathcal{O}_X$. Let $\{v_1, \dots, v_m\}$ be another basis of $X_{\mathcal{A}}$. Then $v_j = \sum_i u_i(u_i|v_j)_{\mathcal{A}}$. For $T \in \mathcal{A}' \cap \mathcal{O}_X$, we have

$$\begin{aligned} \sum_{j=1}^m v_j T v_j^* &= \sum_j \left(\sum_i u_i(u_i|v_j)_{\mathcal{A}} \right) T \left(\sum_k u_k(u_k|v_j)_{\mathcal{A}}^* \right) \\ &= \sum_k \sum_i u_i \left(\sum_j (u_i|v_j)_{\mathcal{A}} (v_j|u_k)_{\mathcal{A}} \right) T u_k^* = \sum_k \sum_i u_i(u_i|u_k)_{\mathcal{A}} T u_k^* \\ &= \sum_k u_k T u_k^* = \sigma(T). \end{aligned}$$

3.2. Lemma. For $T \in \mathcal{A}' \cap \mathcal{O}_X$ and $x_1, \dots, x_m \in X$ we have

$$\sigma^m(T) x_1 \dots x_m = x_1 \dots x_m T$$

and $\sigma^m(T)$ commutes with $\mathcal{K}_{\mathcal{A}}(X^{\otimes m}_{\mathcal{A}})$. In particular $\sigma(T)$ commutes with $\mathcal{A} \subset \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}})$ and σ preserves $\mathcal{A}' \cap \mathcal{O}_X$.

Proof.

$$\begin{aligned} \sigma^m(T) x_1 \dots x_m &= \sum_{i_m} \dots \sum_{i_1} u_{i_m} \dots u_{i_1} T u_{i_1}^* \dots u_{i_m}^* x_1 \dots x_m \\ &= \sum_{i_m} \dots \sum_{i_1} u_{i_m} \dots u_{i_1} u_{i_1}^* \dots u_{i_m}^* x_1 \dots x_m T = x_1 \dots x_m T. \end{aligned}$$

Therefore for $x_1, \dots, x_m, y_1, \dots, y_m \in X$,

$$\begin{aligned} \sigma^m(T) x_1 \dots x_m y_m^* \dots y_1^* &= x_1 \dots x_m T y_m^* \dots y_1^* \\ &= x_1 \dots x_m y_m^* \dots y_1^* \sigma^m(T) \end{aligned}$$

The following Lemma shows that σ acts as a shift operator on the subspaces $\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}}, X^{\otimes s}_{\mathcal{A}})$.

3.3. Lemma. Let

$$\psi : \oplus_{k \in \mathbb{Z}} \mathcal{O}_{\rho}^{(k)} \rightarrow \oplus_{k \in \mathbb{Z}} \mathcal{F}_{\infty}^{(k)}$$

be the *-isomorphism defined in Lemma 2.4. For $T \in \mathcal{A}' \cap \mathcal{K}_{\mathcal{A}}(X^{\otimes m}_{\mathcal{A}})$, we have $\psi^{-1} \sigma \psi(T) = I \otimes T \in \mathcal{K}_{\mathcal{A}}(X \otimes_{\mathcal{A}} X^{\otimes m}_{\mathcal{A}})$

Proof. Since T commutes with the left action of $\mathcal{A}$, $I \otimes T$ makes sense. Let $T = \sum_{x,y} \theta_{x,y}$, where $x = x_1 \otimes \dots \otimes x_m$ and $y = y_1 \otimes \dots \otimes y_m \in X^{\otimes m}$ run over a finite set of simple tensors. Then

$$\begin{aligned} \psi \sigma \psi^{-1}(T) &= \psi \sigma \psi^{-1} \left(\sum_{x,y} \theta_{x,y} \right) = \psi \sigma \left(\sum_{x,y} S_x S_y^* \right) \\ &= \psi \left(\sum_{i=1}^n \sum_{x,y} S_{u_i} S_x S_y^* S_{u_i}^* \right) = \sum_{i=1}^n \sum_{x,y} \theta_{u_i \otimes x, u_i \otimes y} = I \otimes T. \end{aligned}$$

4. THE IDEAL STRUCTURE OF $\mathcal{O}_X$

In this section we study the ideal structure of $\mathcal{O}_X$ under certain assumptions on the bimodule X . In particular, we get a simplicity criterion for $\mathcal{O}_X$.

It was shown in [El], [Ki] that discrete crossed product C^* -algebras are simple if the group action satisfies an outerness property. On the other hand, Cuntz–Krieger algebras $\mathcal{O}_A$ are simple if the defining matrix A is irreducible and non permutation [CK]. More generally, if A satisfies property (I) of [CK] then $\mathcal{O}_A$ is uniquely determined by generators and relations. Furthermore if $\mathcal{A}$ satisfies property (II) (which is stronger than (I)) then the ideal structure of $\mathcal{O}_A$ is explicitly determined in [Cu2]. Matsumoto generalized the simplicity argument to C^* -algebras associated with subshifts [Ma]. Furthermore, Doplicher–Roberts algebras $\mathcal{O}_\rho$ were shown to be simple in the case where ρ is a unitary finite dimensional representation of a compact group of determinant 1.

Our aim is to formulate a criterion to study the ideal structure of $\mathcal{O}_X$ that generalizes the mentioned results.

4.1. Definition. Let $\mathcal{A}$ be a unital C^* -algebra and $X_{\mathcal{A}}$ a finite projective right Hilbert $\mathcal{A}$ -module. Let $\phi : \mathcal{A} \rightarrow \mathcal{K} = \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$ be a unital isometric $*$ -homomorphism. Then a closed ideal J of $\mathcal{A}$ is called X -invariant if $(x|\phi(a)y)_{\mathcal{A}} \in J$ for $x, y \in X$ and $a \in J$.

We note that $J_X := \{a \in \mathcal{A} : (x|\phi(a)y)_{\mathcal{A}} \in J, x, y \in X\}$ is a closed X -invariant ideal of $\mathcal{A}$ containing J , if J is X -invariant. Note that if X is not full and J is the proper ideal generated by inner products then $J_X = \mathcal{A}$. Otherwise, if X is full then J_X is proper if J is. In particular, if J is a maximal proper X -invariant ideal then $J_X = J$. An algebra $\mathcal{A}$ with no proper X -invariant ideal will be called X -simple.

For any closed ideal J of $\mathcal{A}$ we denote by X_J the closed subset of X of elements $x \in X$ for which $(x|x)_{\mathcal{A}} \in J$. If we approximate x with xu_α , with u_α an approximate unit of J , we see that $(y|x)_{\mathcal{A}} \in J$ for all $y \in X$. This shows in particular that $x + y \in X_J$ if $x, y \in X_J$, i.e. X_J is a subspace of X . Furthermore, since $XJ \subseteq X_J$ and $X_J\mathcal{A} \subseteq X_J$, X/X_J is a right Hilbert module over $\mathcal{A}/J$ in a natural way.

4.2. Proposition. If J is a closed X -invariant ideal of $\mathcal{A}$ then $\tilde{\phi} : \mathcal{A}/J \rightarrow \mathcal{L}_{\mathcal{A}/J}(X/X_J)$, $\tilde{\phi}([a])[x] = [\phi(a)x]$ defines a $*$ -homomorphism such that $\ker \tilde{\phi} = J_X/J$. In particular, if $J_X = J$ then $\tilde{\phi}$ is isometric.

Proof. Since X is finite projective and J is X -invariant, $\phi(J)X \subseteq XJ \subseteq X_J$. Furthermore for $x \in X_J$, $a \in \mathcal{A}$, $(\phi(a)x|\phi(a)x) = (x|\phi(a^*a)x) \in J$, thus $\phi(\mathcal{A})X_J \subseteq X_J$. It follows that $\tilde{\phi}$ is a well defined $*$ -homomorphism. The rest of the proof is now clear.

A bimodule X is called (I)-free if for any $k \in \mathbb{N}$ there exists an element $T_k \in \mathcal{A}' \cap \mathcal{L}_{\mathcal{A}}(X^{\otimes p_k}_{\mathcal{A}}, X^{\otimes q_k}_{\mathcal{A}})$ of norm 1 satisfying the following conditions:

$$\mathcal{A} \ni a \mapsto \phi(a)T_k^*T_k \in \mathcal{K}_{\mathcal{A}}(X^{\otimes r_k}_{\mathcal{A}}) \quad (4.1)$$

is completely isometric and

$$\|T_k^* \sigma^k(T_k)\| < 1. \quad (4.2)$$

A bimodule X is called (II) -free if for any closed X -invariant ideal J of $\mathcal{A}$ such that $J_X = J$, X/X_J is (I) -free. If $J = \{0\}$ then $J_X = \{0\}$ since ϕ is faithful, therefore (II) -freeness implies (I) -freeness.

Let $(V, \rho_{\mathcal{A}})$ be a representation of the Hilbert $\mathcal{A}$ -bimodule X in $\mathcal{D}$ with support I . Let $C^*(V(X))$ and $*alg(V(X))$ denote respectively the C^* -subalgebra and the $*$ -subalgebra of $\mathcal{D}$ generated by $V(X)$.

The following Theorem shows that if X is (I) -free, then generators and relations determine $\mathcal{O}_X$ uniquely, whenever $\mathcal{A}$ is faithfully represented. If in addition X is (II) -free then one can determine explicitly the ideal structure of $\mathcal{O}_X$.

4.3. Theorem. *Let X be a full Hilbert module over a C^* -algebra $\mathcal{A}$ and let $\mathcal{D}$ be a unital C^* -algebra, $(V, \rho_{\mathcal{A}})$ a representation of ${}_A X_A$ in $\mathcal{D}$ with support I and let $\varphi : \mathcal{O}_X \rightarrow \mathcal{D}$ be the unique $*$ -homomorphism such that $\varphi(x) = V_x$ and $\varphi(a) = \rho_{\mathcal{A}}(a)$ for $x \in X, a \in \mathcal{A}$ (by the universality of $\mathcal{O}_X$).*

(i) *If $\mathcal{A} \ni a \rightarrow \rho_{\mathcal{A}}(a) \in \mathcal{D}$ is one to one, then the restriction of φ to $\mathcal{O}_X^{(0)}$ is one to one. Furthermore, if X is (I) -free then*

- (1) *$\varphi : \mathcal{O}_X \rightarrow \mathcal{D}$ is one to one, hence ${}^0\mathcal{O}_X \cong *alg(V(X))$,*
- (2) *${}^0\mathcal{O}_X$ has a unique C^* -norm.*

(ii) *If X is (II) -free then*

- (1) *' the maps $\mathcal{J} \mapsto J := \mathcal{J} \cap \mathcal{A}, J \rightarrow \mathcal{J} := cls\{X^r J X^{s*}, r, s = 0, 1, 2, \dots\}$ are inclusion preserving bijective correspondences, inverse of one another, between closed ideals of $\mathcal{O}_X$ and X -invariant ideals J of $\mathcal{A}$ for which $J_X = J$.*
- (2) *' if J is the ideal of $\mathcal{A}$ corresponding to the ideal $\mathcal{J}$ of $\mathcal{O}_X$ as in (1)' then $\mathcal{O}_X/\mathcal{J} = \mathcal{O}_{X/X_J}$.*

Proof. (i): Apply Lemma 2.2 to $X^{\otimes r}$ in place of X , for any r . The restriction of φ to $\mathcal{K}_{\mathcal{A}}(X^{\otimes m} {}_A)$ is isometric, hence the restriction of φ to $\mathcal{O}_X^{(0)}$ is isometric. Let us assume that X is (I) -free.

We claim, without proving it, that the arguments of Lemma 2.2 can be easily generalized to show that for any pair of Hilbert bimodules X, Y and any $x_1, \dots, x_n, y_1, \dots, y_n \in X, T \in \phi(\mathcal{A})' \cap \mathcal{L}_{\mathcal{A}}(Y_A)$,

$$\left\| \left(\sum_{i=1}^n \theta_{x_i, y_i} \otimes 1_Y \right) \circ (1_X \otimes T) \right\| = \left\| ((x_i | x_j)_{\mathcal{A}})^{1/2} ((y_i | y_j)_{\mathcal{A}})^{1/2} diag((T^* T)^{1/2}) \right\|$$

where the right hand side is the norm on $M_n(\phi(\mathcal{A})' \cap \mathcal{L}_{\mathcal{A}}(Y_A))$ and $diag(B)$ is the matrix of $M_n(\mathcal{A})$ with B in the diagonal. It follows that (4.1) can be also read

$$\|\sigma^p(T_n^* T_n) T\| = \|T\|, \quad T \in \mathcal{K}_{\mathcal{A}}(X^{\otimes p} {}_A)$$

Let r_n be sufficiently large so that $T_n^* \sigma^j(T_n) \in \mathcal{L}_{\mathcal{A}}(X^{\otimes r_n} {}_A)$ for $j = 1, \dots, n$. We claim also that replacing r_n by some other r'_n , if necessary, we may find T'_n satisfying the above property and the following property, stronger than (4.2),

$$\|T'_n{}^* \sigma^n(T'_n)\| < \varepsilon.$$

To prove the claim it is enough to assume $n = 1$, since we may replace $T = T_1$ by T_n , r_1 by r_n and X by $X^{\otimes n}$. Choose a natural number q such that $\|T^*\sigma(T)\|^{q+1} < \varepsilon$. Then $T' := T\sigma^{r_1}(T) \dots \sigma^{qr_1}(T)$ satisfies the desired properties. Indeed, using the fact that $T^*\sigma(T) \in \mathcal{L}_{\mathcal{A}}(X^{\otimes r_1}\mathcal{A})$ commutes with $\sigma^{r_1}(T^*)$, $\sigma^{2r_1}(T^*)$, $\dots$, $\sigma^{qr_1}(T^*)$, we have that

$$\begin{aligned} \|T'^*\sigma(T')\| &= \|T^*\sigma(T)\sigma^{r_1}(T^*\sigma(T))\sigma^{2r_1}(T^*\sigma(T)) \dots \sigma^{qr_1}(T^*\sigma(T))\| \\ &\leq \|T^*\sigma(T)\|^{q+1} < \varepsilon. \end{aligned}$$

Moreover, for any $Z \in \mathcal{L}_{\mathcal{A}}(X^{\otimes p}\mathcal{A})$,

$$\|\sigma^p(T'^*T')Z\| = \|Z\sigma^p(T^*T)\sigma^{p+r_1}(T^*T) \dots \sigma^{p+qr_1}(T^*T)\| = \|Z\|.$$

Let now $B = \sum_{j=-n}^n B_j$ be an element of ${}^0\mathcal{O}_X$, with

$$B_j \in \mathcal{L}_{\mathcal{A}}(X^{\otimes p}\mathcal{A}, X^{\otimes p+j}\mathcal{A})$$

and $n \neq 0$. Note that setting

$$B' := \sigma^p(T'_n{}^*)B\sigma^p(T'_n),$$

$$B' = \sum_{j=-n}^n B'_j,$$

where for non negative j ,

$$B'_j = \sigma^p(T'_n{}^*)\sigma^j(T'_n)B_j.$$

It follows that for some p' and all j ,

$$B'_j \in \mathcal{L}_{\mathcal{A}}(X^{\otimes p'}\mathcal{A}, X^{\otimes p'+j}\mathcal{A}).$$

Using the fact that the homogeneous subspaces ${}^0\mathcal{O}_X^{(k)}$ have a unique C^* -norm and that φ is faithful, hence isometric, on them, we deduce, by property (4.1) that

$$\|\varphi(B'_0)\| = \|B'_0\| = \|B_0\|,$$

furthermore,

$$\begin{aligned} \|B'_j\| &< \varepsilon \|B_j\| \quad \text{for } j = -n, n, \\ \|B'_j\| &\leq \|B_j\| \quad \text{for } j = -(n-1), \dots, n-1, \\ \|\varphi(B')\| &\leq \|\varphi(B)\|. \end{aligned}$$

Applying the same computation to B' , replacing T'_n by T'_{n-1} , we deduce, after a finite number of steps, by the arbitrariness of ε , that

$$\|B_0\| = \|\varphi(B_0)\| \leq \|\varphi(B)\|.$$

Thus there exists a conditional expectation $\widehat{E} : \varphi(\mathcal{O}_X) \rightarrow \varphi(\mathcal{O}_X^{(0)})$ such that $\widehat{E}(\varphi(B)) = \varphi(B_0)$. Since $\widehat{E}\varphi = \varphi m_0$, φ is one to one on $\mathcal{O}_X^{(0)}$ and m_0 is faithful, we have that φ is one to one, and (1) follows. On the other hand we have just shown that the inequality $\|B_0\| \leq \|B\|$ holds in any C^* -norm on ${}^0\mathcal{O}_X$, hence the proof of (2) is complete for, by [DR1], ${}^0\mathcal{O}_X$ has a unique C^* -norm with this property.

(ii): Let now assume that X is (II)-free and let $\mathcal{J}$ be a closed ideal of $\mathcal{O}_X$. Then $J = \mathcal{J} \cap \mathcal{A}$ is a closed X -invariant ideal of $\mathcal{A}$ satisfying $J_X = J$. Hence X/X_J is (I)-free. Since $X_J = \mathcal{J} \cap X$, the bimodule X/X_J is isometrically isomorphic to $X/\mathcal{J}$ thought of as a Hilbert bimodule over $\mathcal{A}/J$, thus there is a natural representation of X/X_J in $\mathcal{O}_X/\mathcal{J}$ that is faithful on $\mathcal{A}/J$. We may apply a) to deduce that $\mathcal{O}_{X/X_J}$ is isomorphic to $\mathcal{O}_X/\mathcal{J}$. Let m_k denote the projection onto ${}^0\mathcal{O}_X^{(k)}$. The first part of the proof applied to this representation also shows that if B is an element of $\mathcal{J}$ then $m_0(B)$ is still in $\mathcal{J}$. Now for positive k , for any $x \in X^k$, $x^*m_k(B) = m_0(x^*B)$, thus $x^*m_k(B) \in \mathcal{J}$, hence $m_k(B) \in \mathcal{J}$. Since, by Fourier analysis, any element B of $\mathcal{O}_X$ is the limit in norm of Cesaro sums of $m_k(B)$'s, we deduce that any closed ideal $\mathcal{J}$ is invariant under the canonical action of $\mathbb{T}$, and hence it is generated by the subspaces $\mathcal{J} \cap \mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}}, X^{\otimes s}_{\mathcal{A}}) = X^s J X^{r*}$, therefore the correspondence is one to one. It remains to show that the correspondence is surjective, i.e. that for any X -invariant ideal J of $\mathcal{A}$ satisfying $J_X = J$ we have $\mathcal{J} \cap \mathcal{A} = J$, where $\mathcal{J}$ is the closed ideal of $\mathcal{O}_X$ generated by J . Clearly $\mathcal{J}$ is invariant under the automorphic action of $\mathbb{T}$, hence $J \subseteq \mathcal{J}^{(0)} \cap \mathcal{A}$. Now $\mathcal{J}^{(0)}$ is the inductive limit of the $X^r J X^{r*}$, and $X^r J X^{r*}$ is a closed ideal of $\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})$ that we will denote by $\mathcal{J}_r$. Let π_r be the composition of the inclusion of $\mathcal{A}$ in $\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})$ with the quotient map of $\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})$ in $\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})/\mathcal{J}_r$. Similarly, let π_{∞} be the composition of the inclusion of $\mathcal{A}$ in $\mathcal{O}_X^{(0)}$ with the quotient map of $\mathcal{O}_X^{(0)}$ in $\mathcal{O}_X^{(0)}/\mathcal{J}^{(0)}$. Then $\ker \pi_r = \mathcal{A} \cap \mathcal{J}_r$ and $\ker \pi_{\infty} = \mathcal{A} \cap \mathcal{J}^{(0)}$, hence there are obvious $*$ -monomorphisms

$$\mathcal{A}/\mathcal{A} \cap \mathcal{J}_r \rightarrow \mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})/\mathcal{J}_r$$

$$\mathcal{A}/\mathcal{A} \cap \mathcal{J}^{(0)} \rightarrow \mathcal{O}_X^{(0)}/\mathcal{J}^{(0)}.$$

If b is an element of a C^* -algebra $\mathcal{B}$ and $\mathcal{I}$ is a closed ideal of $\mathcal{B}$, we denote by $[b]_{\mathcal{B}/\mathcal{I}}$ the image of b under the quotient map $\mathcal{B} \rightarrow \mathcal{B}/\mathcal{I}$.

Since a monomorphism between C^* -algebras is isometric, for any $a \in \mathcal{A}$

$$\|[a]_{\mathcal{A}/\mathcal{A} \cap \mathcal{J}^{(0)}}\| = \|[a]_{\mathcal{O}_X^{(0)}/\mathcal{J}^{(0)}}\| = \text{dist}(a, \mathcal{J}^{(0)}) =$$

$$\lim_r \text{dist}(a, \mathcal{J}_r) = \lim_r \|[a]_{\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})/\mathcal{J}_r}\| = \lim_r \|[a]_{\mathcal{A}/\mathcal{A} \cap \mathcal{J}_r}\|.$$

We claim that $J_X = J$ implies that $\mathcal{A} \cap \mathcal{J}_r = J$ for all r , hence adding this fact to the above chain of equalities we deduce that

$$\|[a]_{\mathcal{A}/\mathcal{A} \cap \mathcal{J}^{(0)}}\| = \|[a]_{\mathcal{A}/J}\|.$$

It follows that the natural $*$ -homomorphism $\mathcal{A}/J \rightarrow \mathcal{A}/\mathcal{A} \cap \mathcal{J}^{(0)}$ is injective, i.e. $\mathcal{A} \cap \mathcal{J}^{(0)} = J$. We finally prove the claim. If $a \in X^r J X^{r*} \cap \mathcal{A}$ then for $x, y \in X^r$, $x^* a y \in J$ hence for $x', y' \in X^{r-1}$, $x'^* a y' \in J_X = J$. Iterating the argument we deduce that $a \in J$, and the proof is complete.

4.4. Corollary. *Let $\mathcal{A}$ be a unital C^* -algebra and ${}_A X_{\mathcal{A}}$ a Hilbert $\mathcal{A}$ -bimodule, full and finite projective as a right Hilbert $\mathcal{A}$ -module. Let $\phi : \mathcal{A} \rightarrow \mathcal{K} = \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$ be the defining left action of $\mathcal{A}$ on X . If X is (I) -free and $\mathcal{A}$ is X -simple, then the C^* -algebra $\mathcal{O}_X$ is simple.*

Proof. Let $\mathcal{J} \neq \mathcal{O}_X$ be a closed ideal of $\mathcal{O}_X$. Since $J := \mathcal{J} \cap \mathcal{A}$ is X -invariant and $\mathcal{A}$ is X -simple, we have that $J = 0$. Then the image of $\mathcal{A}$ in $\mathcal{O}_X/\mathcal{J}$ under the quotient map is faithful and all the relations are preserved. Thus the quotient map is one to one by Theorem 4.3 (1), that is $\mathcal{J} = 0$. Therefore $\mathcal{O}_X$ is simple.

Remark 1. Let X be any Hilbert bimodule over $\mathcal{A}$. It has been proved in [DPZ] that X -simplicity of $\mathcal{A}$ is a necessary condition to get simplicity of $\mathcal{O}_X$, but it is not sufficient in general. Similarly to the case of discrete crossed products [El,Ki], one needs to assure furthermore that a certain Connes spectrum naturally associated with X is full [DPZ].

Remark 2. Following more closely the techniques of Elliott [El] and Kishimoto [Ki], it is easy to see that we can replace the assumption of (I) -freeness in Corollary 4.4 by the following condition (i) or a weaker condition (ii).

(i) For any $k \in \mathbb{N}$, any $z \in \mathcal{L}_{\mathcal{A}}(X^{\otimes k}_{\mathcal{A}})$, any finite subset $\{z_1, \dots, z_m\} \subset \mathcal{L}_{\mathcal{A}}(X^{\otimes k}_{\mathcal{A}})$ and any $\epsilon > 0$, there exists $q \in \mathcal{O}_X^{(0)}$ with $\|q\| = 1$ such that $\|qzq\| \geq \|z\| - \epsilon$, $\|qz_i - z_iq\| \leq \epsilon$ for $i = 1, \dots, m$ and $\|q\sigma^j(q)\| \leq \epsilon$ for $j = 1, \dots, k$.

(ii) For any $k \in \mathbb{N}$, any $z \in \mathcal{L}_{\mathcal{A}}(X^{\otimes k}_{\mathcal{A}})$, any finite subset $\{z_1, \dots, z_m\} \subset \mathcal{L}_{\mathcal{A}}(X^{\otimes k}_{\mathcal{A}})$ and any $\epsilon > 0$, there exists $q \in \mathcal{O}_X^{(0)}$ with $\|q\| = 1$ such that $\|qzq\| \geq \|z\| - \epsilon$, $\|qz_i\sigma^j(q)\| \leq \epsilon$ and $\|\sigma^j(q)z_iq\| \leq \epsilon$ for $i = 1, \dots, m$ and $j = 1, \dots, k$.

Remark 3. We can also provide a proof of the simplicity of the C^* -algebra $\mathcal{O}_{\Lambda}$ associated with a certain class of general subshifts studied by Matsumoto [Ma] and in particular $\mathcal{O}_{\beta}$ associated with β -shifts studied in [KMW] by our argument.

5. APPLICATIONS OF THE SIMPLICITY CRITERION

a) Real or Pseudoreal Bimodules

Jones Index theory for C^* -subalgebras was developed in [Wa,KW1] based on the idea that if $\mathcal{A} \subset \mathcal{B}$ is an inclusion of unital C^* -algebras with finite index, $\mathcal{B}$ should be regarded as a finite projective module over $\mathcal{A}$. More specifically, a conditional expectation $E : \mathcal{B} \rightarrow \mathcal{A}$ has finite index if there is a finite subset $\{u_1, \dots, u_n\} \subset \mathcal{B}$, called a basis, such that $x = \sum_{i=1}^n u_i E(u_i^* x)$ for all $x \in \mathcal{B}$. In [Wa], $\{(u_1, u_1^*), \dots, (u_n, u_n^*)\}$ is called a quasi-basis. The index of E is defined by $\text{Index } E = \sum_i u_i u_i^*$. It is shown that $\text{Index } E$ does not depend on the choice of basis and $\text{Index } E$ is in the center of $\mathcal{B}$. We see that $\text{Index } E \geq I$ and $\text{Index } E = I$ if and only if $\mathcal{A} = \mathcal{B}$. We consider $X = \mathcal{B}_{\mathcal{A}}$ as a right Hilbert $\mathcal{A}$ -module with the $\mathcal{A}$ -valued inner product $(x|y)_{\mathcal{A}} = E(x^* y)$. Define a unital isometric $*$ -homomorphism $\phi : \mathcal{A} \rightarrow \mathcal{K} = \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}}) = \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}})$ by $\phi(a)x = ax$ for $a \in \mathcal{A}$ and $x \in X = \mathcal{B}$.

More generally, in [KW1] the authors considered the concept of Hilbert bimodules of finite type. We denote by $d(X)$ the (minimal) dimension of a Hilbert bimodule of finite type.

Let us regard X as an object of the tensor C^* -category $\mathcal{H}_{\mathcal{A}}^b$ with objects Hilbert $\mathcal{A}$ -bimodules and arrows between two objects X and Y the subspace ${}_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}}, Y_{\mathcal{A}})$ of $\mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}}, Y_{\mathcal{A}})$ of operators that commute with the left $\mathcal{A}$ -action. Assume that X has finite intrinsic dimension in $\mathcal{H}_{\mathcal{A}}^b$ in the sense of [LR], i.e. there are arrows $R \in {}_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(\mathcal{A}_{\mathcal{A}}, Y_{\mathcal{A}} \otimes_{\mathcal{A}} X_{\mathcal{A}})$ and $\overline{R} \in {}_{\mathcal{A}}\mathcal{L}_{\mathcal{A}}(\mathcal{A}_{\mathcal{A}}, X_{\mathcal{A}} \otimes_{\mathcal{A}} Y_{\mathcal{A}})$ satisfying the conjugate equations

$$\overline{R}^* \otimes 1_X \circ 1_X \otimes R = 1_X ,$$

$$R^* \otimes 1_Y \circ 1_Y \otimes \overline{R} = 1_Y .$$

X is called real (resp. pseudoreal) if there is a solution of the conjugate equations with $Y = X$ and $\overline{R} = R$ ($\overline{R} = -R$ resp.).

Let $\mathcal{A}$ be a unital C^* -algebra, and let X be a Hilbert bimodule over $\mathcal{A}$ such that $X_{\mathcal{A}}$ is finite projective. It is then straightforward to check that X is real (resp. pseudoreal) if and only if there is an invertible $F : X \rightarrow X$ such that

$$F(axa') = a'^* F(x) a^* ,$$

for $a, a' \in \mathcal{A}$, $x \in X$, and satisfying $F^2 = I$ ($F^2 = -I$ resp.).

The bimodule arising from an inclusion of C^* -algebras $\mathcal{A} \subset \mathcal{B}$ with finite Jones index in the sense of [Wa] is a typical example of real Hilbert $\mathcal{A}$ -bimodule.

5.1. Theorem. *Let X be a Hilbert $\mathcal{A}$ -bimodule, with real or pseudoreal structure defined by an element R such that $\|(R^*R)^{-1}\| < 1$. Then the map $\mathcal{J} \rightarrow \mathcal{J} \cap \mathcal{A}$ is a bijective correspondence between closed ideals of $\mathcal{O}_X$ and X -invariant ideals of $\mathcal{A}$. In particular $\mathcal{O}_X$ is simple if and only if $\mathcal{A}$ is X -simple. Furthermore, if $\mathcal{A}$ is nonnuclear, then $\mathcal{O}_X$ is nonnuclear.*

Proof. By Theorem 4.3 we need to show that every closed X -invariant ideal J of $\mathcal{A}$ satisfies $J_X = J$ and that X is (II) -free.

To prove the first claim it is enough to show that for some $s \in \mathbb{N}$, $X^{\otimes s}$ contains a central element S (i.e. $aS = Sa$, $a \in \mathcal{A}$) satisfying $(S|S)_{\mathcal{A}} = I$, since, then, if $a \in J_X$ we have $a = (S|Sa)_{\mathcal{A}} = (S|\phi(a)S)_{\mathcal{A}} \in J$.

We set $S := R(R^*R)^{-1/2} \in X^{\otimes 2}$. We have that

$$S^* \sigma(S) = (R^*R)^{-1/2} \sigma((R^*R)^{-1/2}) \in \phi(\mathcal{A})' \cap \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}}),$$

hence $\|S^* \sigma(S)\| \leq \|(R^*R)^{-1}\| < 1$. We check that the isometries $S_n := \sigma^{n-1}(S) \dots \sigma(S)S \in \mathcal{A}' \cap X^{2n}$ satisfy the required properties for (I) -freeness. Now (4.1) is a consequence of the fact that S_n is an isometry. We prove (4.2). $S_n^* \sigma^n(S_n) = S_{n-1}^* \sigma^{n-1}(S_{n-1}) \sigma^{n-1}(S^* \sigma(S))$ hence $\|S_n^* \sigma^n(S_n)\| < 1$. We now show that X is (II) -free. Let J be a closed X -invariant ideal of $\mathcal{A}$. For any $n \in \mathbb{N}$ let $\pi_n : X^{\otimes n} \rightarrow (X/X_J)^{\otimes n}$ denote the n -th tensor power of the quotient map $\pi : X \rightarrow X/X_J$. Then the elements $\pi_{2n}(S_{2n}) \in (X/X_J)^{\otimes 2n}$ are still isometries in $(\mathcal{A}/J)' \cap (X/X_J)^{\otimes 2n}$ that satisfy the required properties.

Finally if $\mathcal{A}$ is nonnuclear, then $\mathcal{O}_X$ is nonnuclear, because there exists a conditional expectation $E_{\mathcal{A}}^{\mathcal{O}_X} : \mathcal{O}_X \rightarrow \mathcal{A}$ by Lemma 2.6.

Note that if $d(X)$ is the minimal dimension of X then $d(X)^{-1} \leq \|(R^*R)^{-1}\|$. However, the condition $d(X) > 1$ is not enough to hold the Theorem true

in general (if we do not assume that $\mathcal{A}$ has trivial center) as the following counterexample shows.

Example 1. Let $\mathcal{A}_1 = \mathbb{C} = \mathcal{B}_1$ and $\mathcal{A}_2 = \mathbb{C} \subset \mathcal{B}_2 = \mathbb{C}^2$. Consider conditional expectations $E_1 = id : \mathcal{B}_1 \rightarrow \mathcal{A}_1$ and $E_2 : \mathcal{B}_2 \rightarrow \mathcal{A}_2$ defined by $E_2(x, y) = (\frac{x+y}{2}, \frac{x+y}{2})$. Set $\mathcal{A} = \mathcal{A}_1 \oplus \mathcal{A}_2 \subset \mathcal{B} = \mathcal{B}_1 \oplus \mathcal{B}_2$ and $E = E_1 \oplus E_2 : \mathcal{B} \rightarrow \mathcal{A}$. $X_1 = (\mathcal{B}_1)_{\mathcal{A}_1}$ and $X_2 = (\mathcal{B}_2)_{\mathcal{A}_2}$, $X = \mathcal{B}_{\mathcal{A}}$ with inner products defined by E , and left and right actions of $\mathcal{A}$ defined by multiplication. Since $X = X_1 \oplus X_2$, we have

$$\mathcal{O}_X \cong \mathcal{O}_{X_1} \oplus \mathcal{O}_{X_2} \cong \mathbb{C}(\mathbb{T}) \oplus \mathcal{O}_2$$

Thus it is clear that $\mathcal{O}_X$ is not simple. We have $d(X) > 1$ but there is no bijective correspondence between ideals, because $\mathcal{O}_1$ is isomorphic to $\mathbb{C}(\mathbb{T})$. In fact X is not (II) -free since $X/X_{\mathcal{A}_2} = \mathbb{C}$ is not (I) -free.

However, if we assume that $\mathcal{A}$ is X -simple, the conclusions of the above Theorem hold in the following important particular case.

5.2. Theorem. *Let $\mathcal{A} \subset \mathcal{B}$ be an inclusion of unital C^* -algebras. Let $E : \mathcal{B} \rightarrow \mathcal{A}$ be a conditional expectation with finite index in the sense of [Wa]. Consider $X = \mathcal{B}_{\mathcal{A}}$ as a right Hilbert $\mathcal{A}$ -module with $(x|y)_{\mathcal{A}} = E(x^*y)$ for $x, y \in \mathcal{B}$ and define a left $\mathcal{A}$ -action $\phi : \mathcal{A} \rightarrow \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}})$ by $\phi(a)x = ax$ for $a \in \mathcal{A}$ and $x \in X$. If $\mathcal{A}$ is X -simple and $\text{Index } E \neq I$, then the C^* -algebra $\mathcal{O}_X$ is simple.*

Proof. We show that X is (I) -free, so the result is a consequence of Corollary 4.4. Since $\text{Index } E \neq I$, we have that the Jones projection $e_{\mathcal{A}} \neq I$ and there exists an element $x_0 \in X$ such that $e_{\mathcal{A}}(x_0) \neq x_0$. Since $e_{\mathcal{A}} \in \mathcal{A}' \cap \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}})$, we can define $q_k = e_{\mathcal{A}} \otimes \cdots \otimes e_{\mathcal{A}} \otimes (1 - e_{\mathcal{A}}) \in \mathcal{K}_{\mathcal{A}}(X^{\otimes k+1}_{\mathcal{A}})$ for each $k \in \mathbb{N}$. Then it is easy to see that $q_k \in \mathcal{A}' \cap \mathcal{O}_X$ and for any $m \in \mathbb{N}$ with $1 \leq m \leq k$, we have $q_k \sigma^m(q_k) = 0$. Since

$$\begin{aligned} & (q_k(1 \otimes \cdots \otimes 1 \otimes x_0) \mid 1 \otimes \cdots \otimes 1 \otimes (x_0 - e_{\mathcal{A}}(x_0)))_{\mathcal{A}} \\ &= (1 \otimes \cdots \otimes 1 \otimes (x_0 - e_{\mathcal{A}}(x_0)) \mid 1 \otimes \cdots \otimes 1 \otimes (x_0 - e_{\mathcal{A}}(x_0)))_{\mathcal{A}} \\ &= (x_0 - e_{\mathcal{A}}(x_0) \mid x_0 - e_{\mathcal{A}}(x_0))_{\mathcal{A}} \neq 0 \end{aligned}$$

Therefore $q_k \neq 0$. Let $J = \{a \in \mathcal{A}; ab = aE(b) \text{ for all } b \in \mathcal{B}\}$. Then J is a closed ideal of $\mathcal{A}$ and it is X -invariant. In fact for $a \in J$, $x, y, b \in \mathcal{B}$ we have

$$(x|ay)_{\mathcal{A}}b = E(x^*ay)b = E(x^*aE(y))b = E(x^*)aE(E(y)b) = (x|ay)_{\mathcal{A}}E(b).$$

Thus $J = \{0\}$ since $I \notin J$ and $\mathcal{A}$ is X -simple. Now the $*$ -homomorphism $\mathcal{A} \ni a \rightarrow \phi(a)q_k \in \mathcal{O}_X^{(0)}$ has kernel exactly J , hence it is faithful. This shows that X is (I) -free.

Example 2. Let $\mathcal{A}$ be a unital C^* -algebra, G a finite group and $\alpha : G \rightarrow \text{Aut } \mathcal{A}$ an automorphic action. Consider the crossed product $\mathcal{B} = \mathcal{A} \rtimes_{\alpha} G$ and the usual conditional expectation $E : \mathcal{B} \rightarrow \mathcal{A}$ and let $\lambda_g \in \mathcal{B}$ be the unitary in $\mathcal{B}$ implementing α_g . For the inclusion $\mathcal{A} \subset \mathcal{B}$, let $X = \mathcal{B}_{\mathcal{A}}$ be a right Hilbert module with $\phi : \mathcal{A} \rightarrow \mathcal{K}_{\mathcal{A}}(X_{\mathcal{A}})$ as above. If the order n of G is not equal to one and $\mathcal{A}$ is simple, then $\mathcal{O}_X$ is simple. The C^* -algebra $\mathcal{O}_X$ is uniquely determined by the generators $\mathcal{A}$ and n isometries $\{S_g; g \in G\}$ with the relations $aS_g = S_g\alpha_g^{-1}(a)$ and $\sum_g S_g S_g^* = I$. A concrete realization is given

in $\mathcal{B} \otimes \mathcal{O}_n$. Let $\{T_g; g \in G\}$ be the generators of $\mathcal{O}_n$. Then $\mathcal{O}_X$ is generated by $\{a \otimes I; a \in \mathcal{A}\}$ and $\{S_g = \lambda_g \otimes T_g\}$. If the action α is trivial, then $\mathcal{O}_X$ is isomorphic to the tensor product $\mathcal{A} \otimes \mathcal{O}_n$. Therefore $\mathcal{O}_X$ looks like a twisted tensor product of $\mathcal{A}$ by the Cuntz algebra $\mathcal{O}_n$, that is, " $\mathcal{O}_X \cong \mathcal{A} \otimes_\alpha \mathcal{O}_n$ " by an abuse of notation.

Example 3. Let G be a free group of two generators and H a subgroup of G with $[G : H] = n$. Then H is also a free group of $n + 1$ generators. By [Po] we have an inclusion of simple C^* -algebras $\mathcal{A} = C_r^*(H) \subset \mathcal{B} = C_r^*(G)$ and a conditional expectation $E : \mathcal{B} \rightarrow \mathcal{A}$ such that $E(\sum_{g \in G} x_g \lambda_g) = \sum_{h \in H} x_h \lambda_h$. Put $X = \mathcal{B}_\mathcal{A}$ as above. Then $\mathcal{O}_X$ is simple and nonnuclear by Theorem 5.1.

Example 4. Let $e_1, e_2, \dots$ be a sequence of Jones projections such that $e_i e_{i \pm 1} e_i = (4 \cos^2 \pi/5)^{-1} e_i$ for $i = 1, 2, \dots$. Let $\mathcal{B} = C^*\{1, e_1, e_2, e_3, \dots\} \supset \mathcal{A} = C^*\{1, e_2, e_3, \dots\}$ and $E : \mathcal{B} \rightarrow \mathcal{A}$ a conditional expectation of finite index as in [Wa; Example 3.3.14]. Since $\mathcal{A}$ is simple and $\text{Ind } E = 4 \cos^2 \pi/5$, we have that $\mathcal{O}_X$ is simple by Theorem 5.1.

b) Cuntz–Krieger Bimodules

In this subsection we focus our attention on a class of finite projective Hilbert bimodules over finite direct sum of simple unital C^* -algebras. We start with a couple of easy results that we shall need.

5.3. Proposition. *If X is a full Hilbert C^* -bimodule over a unital C^* -algebra $\mathcal{A}$ then there is a multiplet $\hat{y} = \{y_1, \dots, y_n\}$ in X such that*

$$\sum_i (y_i | y_i)_\mathcal{A} = I .$$

The c. p. map $\Phi_{\hat{y}} : T \in \mathcal{O}_X \rightarrow \sum_i y_i^* T y_i \in \mathcal{O}_X$ satisfies

$$\Phi_{\hat{y}}(A\sigma(B)) = \Phi_{\hat{y}}(A)B , A \in \mathcal{O}_X , B \in \mathcal{A}' \cap \mathcal{O}_X .$$

In particular, $\Phi_{\hat{y}}$ is a left inverse of σ on $\mathcal{A}' \cap \mathcal{O}_X$.

Proof. Since X is full and $\mathcal{A}$ is unital, there are elements z_i and z'_i such that $\|\sum_i (z_i | z'_i)_\mathcal{A} - I\| < 1$. In particular $\sum_i (z_i | z'_i)_\mathcal{A}$ is invertible, so we may assume $\sum_i (z_i | z'_i)_\mathcal{A} = 1$. The matrix $(\theta_{z'_i, z'_j})$ is positive, thus there are $T_{i,k} \in \mathcal{L}_\mathcal{A}(X_\mathcal{A})$ such that $(\theta_{z'_i, z'_j}) = \sum_k T_{i,k} T_{j,k}^*$. It follows that

$$\sum_{i,j,k} (T_{k,i}^* z_i | T_{j,k}^* z_j)_\mathcal{A} = I .$$

The elements $y_k = \sum_j T_{j,k}^* z_j$ satisfy the desired properties.

5.4. Lemma. *If $\mathcal{A}$ is unital and X is a full Hilbert bimodule over $\mathcal{A}$ then*

$$\mathcal{Z}(\mathcal{L}_\mathcal{A}(X_\mathcal{A})) = \{x \in X \rightarrow xa \in X , \quad a \in \mathcal{Z}(\mathcal{A}) \}$$

Proof. Clearly the set at the right hand side is contained in the center of $\mathcal{L}_\mathcal{A}(X_\mathcal{A})$. Let $y_1, \dots, y_n$ be a multiplet in X such that $\sum_i (y_i | y_i)_\mathcal{A} = I$. If $T \in \mathcal{Z}(\mathcal{L}_\mathcal{A}(X_\mathcal{A}))$ then for any $x \in X$,

$$Tx = x \sum_i (y_i | T y_i)_\mathcal{A} .$$

Replacing x by xu , with u unitary in $\mathcal{A}$, shows that $a = \sum_i (y_i | T y_i)_{\mathcal{A}}$ is in the center of $\mathcal{A}$, and the proof is complete.

We denote by $r_n : a \in \mathcal{Z}(\mathcal{A}) \rightarrow r_n(a) \in \mathcal{Z}(\mathcal{L}_{\mathcal{A}}(X^{\otimes n} \mathcal{A}))$ the action of $\mathcal{Z}(\mathcal{A})$ on $X^{\otimes n}$ given by right multiplication. We write r for r_1 .

Let $\mathcal{A}$ be the direct sum of $d > 1$ unital simple C^* -algebras,

$$\mathcal{A} = \mathcal{A}_1 \oplus \cdots \oplus \mathcal{A}_d$$

and let $p_1, \dots, p_d$ denote the minimal central projections of $\mathcal{A}$. We associate to X the matrix $A \in M_d(\{0, 1\})$ by

$$A(i, j) = 0, \quad \text{if } \phi(p_i)r(p_j) = 0$$

,

$$A(i, j) = 1, \quad \text{if } \phi(p_i)r(p_j) \neq 0.$$

Recall that A is called *irreducible* if for any pair of i, j there is $n \in \mathbb{N}$ such that $A^n(i, j) > 0$.

5.5. Proposition. *$\mathcal{A}$ is X -simple if and only if the associated matrix A is irreducible.*

Proof. $\mathcal{A}$ is X -simple if and only if for all j , the smallest X -invariant ideal containing p_j is $\mathcal{A}$, that is to say, for all i there is n such that

$$\{(x|\phi(p_j)yp_i) \mid x, y \in X^{\otimes n}\} \neq 0.$$

In other words, $\phi(p_j)r_n(p_i) \neq 0$. To complete the proof we need to show that $\phi(p_i)r(p_{i_1}) \neq 0, \dots, \phi(p_{i_n})r(p_j) \neq 0$ imply $\phi(p_i)r(p_{i_1}) \dots r_n(p_j) \neq 0$ and hence $\phi(p_i)r_n(p_j) \neq 0$. By induction, assume that $\phi(p_i)r(p_{i_1}) \dots r_{n-1}(p_{i_n}) \neq 0$. The map

$$T \in \mathcal{L}_{\mathcal{A}}(X^{\otimes n} \mathcal{A}) \rightarrow r_n(p_j)T$$

is injective on $r_{n-1}(p_{i_n})\mathcal{L}_{\mathcal{A}}(X^{\otimes n} \mathcal{A})$ so the proof is completed by the previous Lemma.

Following Cuntz and Krieger, we consider the commutative C^* -subalgebra $\mathcal{D}_A$ generated by $r_k(p_i)$, $i = 1, \dots, d$, $k = 0, 1, 2, \dots$.

5.6. Proposition. *The spectrum of $\mathcal{D}_A$ is the compact space*

$$\xi_A = \{(x_k)_{k \in \mathbb{N}}, x_k = 1, \dots, d, A(x_k, x_{k+1}) = 1\}.$$

Each projection p_i corresponds to the characteristic function of

$$Z(i) = \{(x_k) \in \xi_A, x_1 = i\}.$$

Furthermore the endomorphism σ acts on ξ_A as the one-sided shift on the left.

Proof. The spectrum of $\mathcal{D}_A$ is given by

$$\{(x_k)_{k \in \mathbb{N}}, x_k = 1, \dots, d, p_{x_1}r(p_{x_2}) \dots r_n(p_{x_{n+1}}) \neq 0, n \in \mathbb{N}\},$$

so the arguments of the previous Proposition complete the proof.

5.7. Theorem. *Let $\mathcal{A}$ be the direct sum of $d > 1$ unital simple C^* -algebras as above. Let X be a finite projective Hilbert bimodule over $\mathcal{A}$. If the associated matrix A satisfies condition (I) of [CK] then X is (I)-free. In particular, if A is irreducible and non permutation then $\mathcal{O}_X$ is a simple C^* -algebra.*

Proof. We can apply the arguments of [CK; Lemmas 2.6, 2.7] to $\mathcal{L}_{\mathcal{A}}(X^{\otimes k}_{\mathcal{A}})$ in place of $\mathcal{F}_k$, and to a basis $\{x_i\}$ of X in place of the partial isometries S_i and deduce that for all $k, r \in \mathbb{N}$ there is a projection $q \in \mathcal{D}_A$ such that

$$\|\sigma^r(q)T\| = \|T\|, \quad T \in \mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}}),$$

$$q\sigma^j(q) = 0, \quad 0 < |j| < k.$$

Approximating q by a projection in some $\mathcal{Z}(\mathcal{L}_{\mathcal{A}}(X^{\otimes p}_{\mathcal{A}}))$ gives an element with the desired properties.

6. BIMODULES OVER PURELY INFINITE C^* -ALGEBRAS

In this section we present a class of examples of purely infinite simple C^* -algebras of the form $\mathcal{O}_X$.

6.1. Theorem. *Let X be a finite projective right Hilbert bimodule over a unital purely infinite simple C^* -algebra $\mathcal{A}$. If X is (I)-free, then $\mathcal{O}_X$ is simple and purely infinite.*

Proof. Note that if Y is a finite projective Hilbert module over $\mathcal{A}$ then $\mathcal{L}_{\mathcal{A}}(Y_{\mathcal{A}})$ is isomorphic to a full corner of some $M_d(\mathcal{A})$, so it is simple and purely infinite. Now $\mathcal{O}_X^{(0)}$ is simple and purely infinite since it is the inductive limit of $\mathcal{L}_{\mathcal{A}}(X^{\otimes r}_{\mathcal{A}})$.

Let T be a non zero positive element in $\mathcal{O}_X$. Since $m_0(T)$ is a non zero positive element in the purely infinite C^* -algebra $\mathcal{O}_X^{(0)}$, there exists $W \in \mathcal{O}_X^{(0)}$ such that $W^*m_0(T)W = 1$. Put $S = W^*TW$. Then $m_0(S) = 1$. Choose $0 < \varepsilon < 1/5$. Then there exists $B \in {}^0\mathcal{O}_X$ with $\|S - B\| < \varepsilon$. Put $B_0 = m_0(B) \in {}^0\mathcal{O}_X^{(0)}$. Then $\|1 - B_0\| = \|m_0(S - B)\| < \varepsilon$. Let $C = B - B_0 + I \in {}^0\mathcal{O}_X$. Then $\|C - B\| < \varepsilon$ and $\|S - C\| < 2\varepsilon$. Consider the expansion $C = \sum_{j=-k}^k C_j \in {}^0\mathcal{O}_X$, with $C_j \in \mathcal{L}_{\mathcal{A}}(X^{\otimes p}_{\mathcal{A}}, X^{\otimes p+j}_{\mathcal{A}})$ for some p . Note that $C_0 = m_0(C) = I$. By the proof of Theorem 4.3, considering the product of operators that appear in the iterating argument, (I)-freeness provides an element $R \in \mathcal{A}' \cap {}^0\mathcal{O}_X$ satisfying $R^*R \in \mathcal{O}_X^{(0)}$, $\|R^*CR - R^*C_0R\| < \varepsilon$ and $\|R^*R\| = \|R^*C_0R\| = \|C_0\| = 1$. Then

$$\|R^*SR - R^*R\| \leq \|R^*(S - C)R\| + \|R^*(C - C_0)R\| < 3\varepsilon$$

Since R^*R has norm one, there exists $V \in \mathcal{O}_X^{(0)}$ such that $V^*R^*RV = I$ and $\|V\| < 1 + \varepsilon$. Then we have

$$\|V^*R^*SRV - I\| = \|V^*R^*SRV - V^*R^*RV\| < 3\varepsilon(1 + \varepsilon)^2 < 108/125 < 1$$

Hence V^*R^*SRV is positive invertible and there exists $Y \in \mathcal{O}_X$ such that $Y^*TY = 1$. Therefore $\mathcal{O}_X$ is purely infinite.

Remark 1. In the above Theorem, instead of assuming $\mathcal{A}$ is purely infinite, we may assume that $\mathcal{A}$ has real rank zero and for any natural number r and

for any non zero projection $P \in \mathcal{L}_{\mathcal{A}}(X_{\mathcal{A}}^{\otimes r})$ there exists $x \in X^{\otimes r}$ with $Px = x$ and $(x|x)_{\mathcal{A}} = 1$. Just note the fact that $x^*x = I$ and

$$I \geq x^*Px \geq x^*\theta_{x,x}x = x^*xx^*x = I$$

Remark 2. If we know a priori that $\mathcal{O}_X$ is simple then we can also conclude, without assuming that X is (I) -free, that $\mathcal{O}_X$ is purely infinite as follows. By [Pe; Theorems 8.10.4; 8.10.10] $\mathcal{O}_X^{(0)}$ has the trivial relative commutant in $\mathcal{O}_X$. Let now x be the image in $\mathcal{O}_X$ of a non zero element in X . By the pure infiniteness of $\mathcal{A}$ there is $a \in \mathcal{A}$ such that $a^*x^*xa = I$, so replacing x by xa , we may assume that x is an isometry. Note that $\mathcal{O}_X$ is generated by $\mathcal{O}_X^{(0)}$ and x . Now the triviality of the relative commutant implies that all the powers of the endomorphism ρ implemented by x on $\mathcal{O}_X^0$ are outer. If x is unitary then $\mathcal{O}_X$ is the crossed product of $\mathcal{O}_X^0$ by the automorphism ρ so it is simple [Ki] and purely infinite [JKO; Theorem 3.2]. If x is a proper isometry, then $\mathcal{O}_X$ is the crossed product of a simple purely infinite C^* -algebra by a proper corner endomorphism, such that all powers are outer so by [JKO; Theorem 2.1], $\mathcal{O}_X$ is simple and purely infinite.

Acknowledgment Part of this paper was written during a visit of C.P. to the Matematisk Institut of Kobenhavn. She wishes to thank Ryszard Nest, Hiroyuki Osaka, Gert K. Pedersen, Carl Winsløw for fruitful discussions and for warm hospitality.

REFERENCES

- [AEE] B. Abadie, S. Eilers and R. Exel, *Morita equivalence for crossed products by Hilbert C^* -bimodules*, Preprint (1995).
- [BMS] L. G. Brown, J. Mingo and N. Shen, *Quasi-multipliers and embeddings of Hilbert C^* -bimodules*, Canad. J. Math., **46**, No 6 (1994), 1150–1174.
- [Co1] A. Connes: *Une classification des facteurs de type III*, Ann. Scient. Èc. Norm. Sup., **6** (1973), 133–252.
- [Co2] A. Connes: *Outer conjugacy classes of automorphisms of factors*, Ann. Scient. Èc. Norm. Sup., **8** (1975), 383–420.
- [CT] A. Connes and M. Takesaki: *The flow of weights on factors of type III*, Tohoku Math. J., **29** (1977), 473–575.
- [Cu] J. Cuntz, *Simple C^* -algebras generated by isometries*, Comm. Math. Phys., **57** (1977), 173–185.
- [CK] J. Cuntz and W. Krieger, *A class of C^* -algebras and topological Markov chains*, Invent. Math., **56** (1980), 251–268.
- [Cu2] J. Cuntz, *A class of C^* -algebras and topological Markov chains II: Reducible chains and the Ext-functor for C^* -algebras*, Invent. Math., **63** (1981), 25–40.
- [DR1] S. Doplicher and J. E. Roberts, *Duals of compact Lie groups realized in the Cuntz algebras and their actions on C^* -algebras*, J. Funct. Anal., **74** (1987), 96–120.
- [DR2] S. Doplicher and J. E. Roberts, *Endomorphisms of C^* -algebras, crossed products and duality for compact groups*, Ann. Math., **130** (1989), 75–119.
- [DR3] S. Doplicher and J. E. Roberts, *A new duality theory for compact groups*, Invent. Math., **98** (1989), 157–218.

- [DPZ] S. Doplicher, C. Pinzari and R. Zuccante, *On the C^* -algebra generated by a Hilbert bimodule*, preprint.
- [El] G. Elliott, *Some simple C^* -algebras constructed as crossed products with discrete outer automorphism groups*, Publ. RIMS, Kyoto Univ., **16** (1980), 299–311.
- [Ex] R. Exel, *Circle actions on C^* -algebras, partial automorphisms and a generalized Pimsner–Voiculescu exact sequence*, J. Funct. Anal., **122** (1994), 361–401.
- [GHJ] F. M. Goodman, P. de la Harpe and V. F. R. Jones, *Coxeter graphs and towers of algebras*, Springer–Verlag, New York, (1989).
- [I1] M. Izumi, *Subalgebras of infinite C^* -algebras with finite Watatani indices. I. Cuntz algebras*, Comm. Math. Phys., **155** (1993), 157–182.
- [I2] M. Izumi, *Subalgebras of infinite C^* -algebras with finite Watatani indices. II. Cuntz–Krieger algebras*, Preprint.
- [I3] M. Izumi, *Index Theory of Simple C^* -algebras*, Talk given at the Fields Institute, 1995.
- [JKO] J. A. Jeong, K. Kodaka and H. Osaka, *Purely infinite simple C^* -crossed products I*, Canad. Math. Bull. Vol., to appear.
- [Jo] V. F. R. Jones, *Index for subfactors*, Invent. Math., **72** (1983), 1–25.
- [KW1] T. Kajiwara and Y. Watatani, *Jones index theory by Hilbert C^* -bimodules and K -theory*, preprint (1996).
- [KW2] T. Kajiwara and Y. Watatani, *Crossed products of Hilbert C^* -bimodules by countable discrete groups*, to appear in Proc. Amer. Math. Soc.
- [KWP] T. Kajiwara, Y. Watatani and C. Pinzari, *Hilbert C^* -bimodules and countably generated Cuntz–Krieger algebras*, in preparation.
- [Ka] Y. Katayama, *Generalized Cuntz algebras $\mathcal{O}_N^M$* , RIMS Kokyuroku, **858** (1994), 131–151.
- [KMW] Y. Katayama, K. Matsumoto and Y. Watatani, *Simple C^* algebras arising from β -expansion of real numbers*, Preprint (1996)
- [Ki] A. Kishimoto, *Outer automorphisms and reduced crossed products of simple C^* -algebras*, Comm. Math. Phys., **81** (1981), 429–435.
- [LR] R. Longo and J. E. Roberts, *A theory of dimension*, to appear in K -theory.
- [Ma] K. Matsumoto, *On C^* -algebras associated with subshifts*, to appear in Internat. J. Math.
- [MS] P. S. Muhly and B. Solel, *On the simplicity of some Cuntz–Pimsner algebras*, preprint (1996).
- [Ol] D. Olesen: *Inner automorphisms of simple C^* -algebras*, Comm. Math. Phys., **44** (1975), 175–190.
- [OP1] D. Olesen and G. K. Pedersen: *Applications of the Connes spectrum to C^* -dynamical systems*, J. Funct. Anal., **30** (1978), 179–197.
- [OP2] D. Olesen and G. K. Pedersen: *Applications of the Connes spectrum to C^* -dynamical systems III*, J. Funct. Anal., **45** (1982), 357–390 .
- [Pe] G. K. Pedersen, *C^* -algebras and their automorphism groups*, Academic Press, New York (1990).
- [PP] M. Pimsner and S. Popa, *Entropy and index for subfactors*, Ann. Sci. Ec. Norm. Sup., **19** (1985), 57–106.
- [Pi] M. Pimsner, *A class of C^* -algebras generalizing both Cuntz–Krieger algebras and Crossed products by $\mathbb{Z}$* , preprint.

- [P] C. Pinzari, *The ideal structure of Cuntz–Krieger–Pimsner algebras and Cuntz–Krieger algebras over infinite matrices*, to appear in the proceedings of the Rome conference “Operator algebras and Quantum Field Theory”, July 1996.
- [Po] R. Powers, *Simplicity of C^* -algebra associated with the free group on two generators*, Duke Math. J., **42** (1975), 151–156.
- [Ri] M. Rieffel, *Induced representations of C^* -algebras*, Adv. Math., **13** (1974), 176–257.
- [Ro] J. E. Roberts, *Cross products of von Neumann algebras by group duals*, Symp. Math., **20** (1976), 335–363.
- [RØ1] M. Rørdam, *Classification of Cuntz–Krieger algebras*, K-theory **9** (1995), 31–58.
- [RØ2] M. Rørdam, *Classification of certain infinite simple C^* -algebras*, J. Funct. Anal., **131** (1995), 415–458.
- [Wa] Y. Watatani, *Index for C^* -subalgebras*, Memoir Amer. Math. Soc., **424** (1990).